

Systems of sets of lengths: transfer Krull monoids versus weakly Krull monoids*

Alfred Geroldinger and Wolfgang A. Schmid and Qinghai Zhong

Abstract Transfer Krull monoids are monoids which allow a weak transfer homomorphism to a commutative Krull monoid, and hence the system of sets of lengths of a transfer Krull monoid coincides with that of the associated commutative Krull monoid. We unveil a couple of new features of the system of sets of lengths of transfer Krull monoids over finite abelian groups G , and we provide a complete description of the system for all groups G having Davenport constant $D(G) = 5$ (these are the smallest groups for which no such descriptions were known so far). Under reasonable algebraic finiteness assumptions, sets of lengths of transfer Krull monoids and of weakly Krull monoids satisfy the Structure Theorem for Sets of Lengths. In spite of this common feature we demonstrate that systems of sets of lengths for a variety of classes of weakly Krull monoids are different from the system of sets of lengths of any transfer Krull monoid.

Key words: transfer Krull monoids, weakly Krull monoids, sets of lengths, zero-sum sequences

Mathematics Subject Classification (2010): 11B30, 11R27, 13A05, 13F05, 16H10, 16U30, 20M13

A. Geroldinger (corresponding author) and Q. Zhong

University of Graz, Institute for Mathematics, Heinrichstraße 36, 8010 Graz, Austria, e-mail: alfred.geroldinger@uni-graz.at, qinghai.zhong@uni-graz.at

W.A. Schmid

Laboratoire Analyse, Géométrie et Applications (LAGA, UMR 7539), COMUE Université Paris Lumières, Université Paris 8, CNRS, 93526 Saint-Denis cedex, France

Université Paris 13, Sorbonne Paris Cité, LAGA, CNRS, UMR 7539, Université Paris 8, F-93430, Villetaneuse, France e-mail: schmid@math.univ-paris13.fr

* This work was supported by the Austrian Science Fund FWF, Project Number P28864-N35, and the ANR project Caesar, project number ANR-12-BS01-0011

1 Introduction

By an atomic monoid we mean a cancellative semigroup with unit element such that every nonunit can be written as a finite product of irreducible elements. Let H be an atomic monoid. If $a \in H$ is a nonunit and $a = u_1 \cdots u_k$ is a factorization of a into k irreducible elements, then k is called a factorization length and the set $L(a) \subset \mathbb{N}$ of all possible factorization lengths is called the set of lengths of a . Then $\mathcal{L}(H) = \{L(a) \mid a \in H\}$ is the system of sets of lengths of H . Under a variety of noetherian conditions on H (e.g., H is the monoid of nonzero elements of a commutative noetherian domain) all sets of lengths are finite. Furthermore, if there is some element $a \in H$ with $|L(a)| > 1$, then $|L(a^N)| > N$ for all $N \in \mathbb{N}$. Sets of lengths (together with invariants controlling their structure, such as elasticities and sets of distances) are a well-studied means of describing the arithmetic structure of monoids ([20, 11]).

Let H be a transfer Krull monoid. Then, by definition, there is a weak transfer homomorphism $\theta: H \rightarrow \mathcal{B}(G_0)$, where $\mathcal{B}(G_0)$ denotes the monoid of zero-sum sequences over a subset G_0 of an abelian group, and hence $\mathcal{L}(H) = \mathcal{L}(\mathcal{B}(G_0))$. A special emphasis has always been on the case where G_0 is a finite abelian group. Thus let G be a finite abelian group and we use the abbreviation $\mathcal{L}(G) = \mathcal{L}(\mathcal{B}(G))$. It is well-known that sets of lengths in $\mathcal{L}(G)$ are highly structured (Proposition 3.2), and the standing conjecture is that the system $\mathcal{L}(G)$ is characteristic for the group G . More precisely, if G' is a finite abelian group such that $\mathcal{L}(G) = \mathcal{L}(G')$, then G and G' are isomorphic (apart from two well-known trivial pairings; see Conjecture 3.4). This conjecture holds true, among others, for groups G having rank at most two, and its proof uses deep results from additive combinatorics which are not available for general groups. Thus there is a need for studying $\mathcal{L}(G)$ with a new approach. In Section 3, we unveil a couple of properties of the system $\mathcal{L}(G)$ which are first steps on a new way towards Conjecture 3.4.

In spite of all abstract work on systems $\mathcal{L}(G)$, they have been written down explicitly only for groups G having Davenport constant $D(G) \leq 4$, and this is not difficult to do (recall that a group G has Davenport constant $D(G) \leq 4$ if and only if either $|G| \leq 4$ or G is an elementary 2-group of rank three). In Section 4 we determine the systems $\mathcal{L}(G)$ for all groups G having Davenport constant $D(G) = 5$.

Commutative Krull monoids are the classic examples of transfer Krull monoids. In recent years a wide range of monoids and domains has been found which are transfer Krull but which are not commutative Krull monoids. Thus the question arose which monoids H have systems $\mathcal{L}(H)$ which are different from systems of sets of lengths of transfer Krull monoids. Commutative v -noetherian weakly Krull monoids and domains are the best investigated class of monoids beyond commutative Krull monoids (numerical monoids as well as one-dimensional noetherian domains are v -noetherian weakly Krull). Clearly, weakly Krull monoids can be half-factorial and half-factorial monoids are transfer Krull monoids. Similarly, it can happen both for weakly Krull monoids as well as for transfer Krull monoids that all sets of lengths are arithmetical progressions with difference 1. Apart from such

extremal cases, we show in Section 5 that systems of sets of lengths of a variety of classes of weakly Krull monoids are different from the system of sets of lengths of any transfer Krull monoid.

2 Background on sets of lengths

We denote by \mathbb{N} the set of positive integers, and for real numbers $a, b \in \mathbb{R}$, we denote by $[a, b] = \{x \in \mathbb{Z} \mid a \leq x \leq b\}$ the discrete interval between a and b , and by an interval we always mean a finite discrete interval of integers.

Let $A, B \subset \mathbb{Z}$ be subsets of the integers. Then $A + B = \{a + b \mid a \in A, b \in B\}$ is the sumset of A and B . We set $-A = \{-a \mid a \in A\}$ and for an integer $m \in \mathbb{Z}$, $m + A = \{m\} + A$ is the shift of A by m . For $m \in \mathbb{N}$, we denote by $mA = A + \dots + A$ the m -fold subset of A and by $m \cdot A = \{ma \mid a \in A\}$ the dilation of A by m . If $A \subset \mathbb{N}$, we denote by $\rho(A) = \sup A / \min A \in \mathbb{Q}_{\geq 1} \cup \{\infty\}$ the elasticity of A and we set $\rho(\{0\}) = 1$. A positive integer $d \in \mathbb{N}$ is called a distance of A if there are $a, b \in A$ with $b - a = d$ and the interval $[a, b]$ contains no further elements of A . We denote by $\Delta(A)$ the set of distances of A . Clearly, $\Delta(A) = \emptyset$ if and only if $|A| \leq 1$, and A is an arithmetical progression if and only if $|\Delta(A)| \leq 1$.

Let G be an additive abelian group. An (ordered) family $(e_i)_{i \in I}$ of elements of G is said to be *independent* if $e_i \neq 0$ for all $i \in I$ and, for every family $(m_i)_{i \in I} \in \mathbb{Z}^{(I)}$,

$$\sum_{i \in I} m_i e_i = 0 \quad \text{implies} \quad m_i e_i = 0 \quad \text{for all } i \in I.$$

A family $(e_i)_{i \in I}$ is called a *basis* for G if $e_i \neq 0$ for all $i \in I$ and $G = \bigoplus_{i \in I} \langle e_i \rangle$. A subset $G_0 \subset G$ is said to be independent if the tuple $(g)_{g \in G_0}$ is independent. For every prime $p \in \mathbb{P}$, we denote by $r_p(G)$ the p -rank of G .

Sets of Lengths. We say that a semigroup S is cancellative if for all elements $a, b, c \in S$, the equation $ab = ac$ implies $b = c$ and the equation $ba = ca$ implies $b = c$. Throughout this manuscript, a monoid means a cancellative semigroup with unit element, and we will use multiplicative notation.

Let H be a monoid. An element $a \in H$ is said to be invertible if there exists an element $a' \in H$ such that $aa' = a'a = 1$. The set of invertible elements of H will be denoted by H^\times , and we say that H is reduced if $H^\times = \{1\}$. For a set P , we denote by $\mathcal{F}(P)$ the free abelian monoid with basis P . Then every $a \in \mathcal{F}(P)$ has a unique representation in the form

$$a = \prod_{p \in P} p^{\nu_p(a)},$$

where $\nu_p: \mathcal{F}(P) \rightarrow \mathbb{N}_0$ denotes the p -adic exponent.

An element $a \in H$ is called *irreducible* (or an *atom*) if $a \notin H^\times$ and if, for all $u, v \in H$, $a = uv$ implies that $u \in H^\times$ or $v \in H^\times$. We denote by $\mathcal{A}(H)$ the set of atoms of H . The monoid H is said to be *atomic* if every $a \in H \setminus H^\times$ is a product of finitely

many atoms of H . If $a \in H$ and $a = u_1 \cdots u_k$, where $k \in \mathbb{N}$ and $u_1, \dots, u_k \in \mathcal{A}(H)$, then we say that k is the *length* of the factorization. For $a \in H \setminus H^\times$, we call

$$\mathsf{L}_H(a) = \mathsf{L}(a) = \{k \in \mathbb{N} \mid a \text{ has a factorization of length } k\} \subset \mathbb{N}$$

the *set of lengths* of a . For convenience, we set $\mathsf{L}(a) = \{0\}$ for all $a \in H^\times$. By definition, H is atomic if and only if $\mathsf{L}(a) \neq \emptyset$ for all $a \in H$. Furthermore, $\mathsf{L}(a) = \{1\}$ if and only if $a \in \mathcal{A}(H)$ if and only if $1 \in \mathsf{L}(a)$. If $a, b \in H$, then $\mathsf{L}(a) + \mathsf{L}(b) \subset \mathsf{L}(ab)$. We call

$$\mathcal{L}(H) = \{\mathsf{L}(a) \mid a \in H\}$$

the *system of sets of lengths* of H . We say that H is *half-factorial* if $|L| = 1$ for every $L \in \mathcal{L}(H)$. If H is atomic, then H is either half-factorial or for every $N \in \mathbb{N}$ there is an element $a_N \in H$ such that $|\mathsf{L}(a_N)| > N$ ([17, Lemma 2.1]). We say that H is a BF-monoid if it is atomic and all sets of lengths are finite. Let

$$\Delta(H) = \bigcup_{L \in \mathcal{L}(H)} \Delta(L) \subset \mathbb{N}$$

denote the *set of distances* of H , and if $\Delta(H) \neq \emptyset$, then $\min \Delta(H) = \gcd \Delta(H)$. We denote by $\Delta_1(H)$ the set of all $d \in \mathbb{N}$ with the following property:

For every $k \in \mathbb{N}$ there exists an $L \in \mathcal{L}(H)$ of the form $L = L' \cup \{y, y+d, \dots, y+kd\} \cup L''$ where $y \in \mathbb{N}$ and $L', L'' \subset \mathbb{Z}$ with $\max L' < y$ and $y+kd < \min L''$.

By definition, $\Delta_1(H)$ is a subset of $\Delta(H)$. For every $k \in \mathbb{N}$ we define the k th *elasticity* of H . If $H = H^\times$, then we set $\rho_k(H) = k$, and if $H \neq H^\times$, then

$$\rho_k(H) = \sup \{\sup L \mid k \in L \in \mathcal{L}(H)\} \in \mathbb{N} \cup \{\infty\}.$$

The invariant

$$\rho(H) = \sup \{\rho(L) \mid L \in \mathcal{L}(H)\} = \lim_{k \rightarrow \infty} \frac{\rho_k(H)}{k} \in \mathbb{R}_{\geq 1} \cup \{\infty\}$$

is called the *elasticity* of H (see [17, Proposition 2.4]). Sets of lengths of all monoids, which are in the focus of the present paper, are highly structured (see Proposition 3.2 and Theorems 5.5 – 5.8). To summarize the relevant concepts, let $d \in \mathbb{N}$, $M \in \mathbb{N}_0$ and $\{0, d\} \subset \mathcal{D} \subset [0, d]$. A subset $L \subset \mathbb{Z}$ is called an *almost arithmetical multiprogression* (AAMP for short) with *difference* d , *period* \mathcal{D} , and *bound* M , if

$$L = y + (L' \cup L^* \cup L'') \subset y + \mathcal{D} + d\mathbb{Z}$$

where $y \in \mathbb{Z}$ is a shift parameter,

- L^* is finite nonempty with $\min L^* = 0$ and $L^* = (\mathcal{D} + d\mathbb{Z}) \cap [0, \max L^*]$, and
- $L' \subset [-M, -1]$ and $L'' \subset \max L^* + [1, M]$.

We say that *the Structure Theorem for Sets of Lengths* holds for a monoid H if H is atomic and there exist some $M \in \mathbb{N}_0$ and a finite nonempty set $\Delta \subset \mathbb{N}$ such that every $L \in \mathcal{L}(H)$ is an AAMP with some difference $d \in \Delta$ and bound M .

Monoids of zero-sum sequences. We discuss a monoid having a combinatorial flavor whose universal role in the study of sets of lengths will become evident at the beginning of the next section. Let G be an additive abelian group and $G_0 \subset G$ a subset. Then $\langle G_0 \rangle$ denotes the subgroup generated by G_0 , and we set $G_0^\bullet = G_0 \setminus \{0\}$. In additive combinatorics, a *sequence* (over G_0) means a finite sequence of terms from G_0 where repetition is allowed and the order of the elements is disregarded, and (as usual) we consider sequences as elements of the free abelian monoid with basis G_0 . Let

$$S = g_1 \cdot \dots \cdot g_\ell = \prod_{g \in G_0} g^{v_g(S)} \in \mathcal{F}(G_0)$$

be a sequence over G_0 . We set $-S = (-g_1) \cdot \dots \cdot (-g_\ell)$, and we call

- $\text{supp}(S) = \{g \in G \mid v_g(S) > 0\} \subset G$ the *support* of S ,
- $|S| = \ell = \sum_{g \in G} v_g(S) \in \mathbb{N}_0$ the *length* of S ,
- $\sigma(S) = \sum_{i=1}^l g_i$ the *sum* of S ,
- $\Sigma(S) = \left\{ \sum_{i \in I} g_i \mid \emptyset \neq I \subset [1, \ell] \right\}$ the *set of subsequence sums* of S ,
- $k(S) = \sum_{i=1}^l \frac{1}{\text{ord}(g_i)}$ the *cross number* of S .

The sequence S is said to be

- *zero-sum free* if $0 \notin \Sigma(S)$,
- *a zero-sum sequence* if $\sigma(S) = 0$,
- *a minimal zero-sum sequence* if it is a nontrivial zero-sum sequence and every proper subsequence is zero-sum free.

The set of zero-sum sequences $\mathcal{B}(G_0) = \{S \in \mathcal{F}(G_0) \mid \sigma(S) = 0\} \subset \mathcal{F}(G_0)$ is a submonoid, and the set of minimal zero-sum sequences is the set of atoms of $\mathcal{B}(G_0)$. For any arithmetical invariant $*(H)$ defined for a monoid H , we write $*(G_0)$ instead of $*(\mathcal{B}(G_0))$. In particular, $\mathcal{A}(G_0) = \mathcal{A}(\mathcal{B}(G_0))$ is the set of atoms of $\mathcal{B}(G_0)$, $\mathcal{L}(G_0) = \mathcal{L}(B(G_0))$ is the system of sets of lengths of $\mathcal{B}(G_0)$, and so on. Furthermore, we say that G_0 is half-factorial if the monoid $\mathcal{B}(G_0)$ is half-factorial. We denote by

$$D(G_0) = \sup\{|S| \mid S \in \mathcal{A}(G_0)\} \in \mathbb{N}_0 \cup \{\infty\}$$

the *Davenport constant* of G_0 . If G_0 is finite, then $D(G_0)$ is finite. Suppose that G is finite, say $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$, with $r \in \mathbb{N}_0$, $1 < n_1 \mid \dots \mid n_r$, then $r = r(G)$ is the rank of G , and we have

$$1 + \sum_{i=1}^r (n_i - 1) \leq D(G) \leq |G|. \quad (1)$$

If G is a p -group or $r(G) \leq 2$, then $1 + \sum_{i=1}^r (n_i - 1) = D(G)$. Suppose that $|G| \geq 3$. We will use that $\Delta(G)$ is an interval with $\min \Delta(G) = 1$ ([26]), and that, for all $k \in \mathbb{N}$,

$$\begin{aligned}
\rho_{2k}(G) &= kD(G), \\
kD(G) + 1 \leq \rho_{2k+1}(G) &\leq kD(G) + \lfloor D(G)/2 \rfloor, \quad \text{and} \\
\rho(G) &= D(G)/2,
\end{aligned} \tag{2}$$

([20, Section 6.3]).

3 Sets of lengths of transfer Krull monoids

Weak transfer homomorphisms play a critical role in factorization theory, in particular in all studies of sets of lengths. We refer to [20] for a detailed presentation of transfer homomorphisms in the commutative setting. Weak transfer homomorphisms (as defined below) were introduced in [5, Definition 2.1] and transfer Krull monoids were introduced in [17].

Definition 3.1. Let H be a monoid.

1. A monoid homomorphism $\theta: H \rightarrow B$ to an atomic monoid B is called a *weak transfer homomorphism* if it has the following two properties:

(T1) $B = B^\times \theta(H)B^\times$ and $\theta^{-1}(B^\times) = H^\times$.

(WT2) If $a \in H$, $n \in \mathbb{N}$, $v_1, \dots, v_n \in \mathcal{A}(B)$ and $\theta(a) = v_1 \cdot \dots \cdot v_n$, then there exist $u_1, \dots, u_n \in \mathcal{A}(H)$ and a permutation $\tau \in \mathfrak{S}_n$ such that $a = u_1 \cdot \dots \cdot u_n$ and $\theta(u_i) \in B^\times v_{\tau(i)}B^\times$ for each $i \in [1, n]$.

2. H is said to be a *transfer Krull monoid* (over G_0) if there exists a weak transfer homomorphism $\theta: H \rightarrow \mathcal{B}(G_0)$ for a subset G_0 of an abelian group G . If G_0 is finite, then we say that H is a *transfer Krull monoid of finite type*.

If R is a domain and R^\bullet its monoid of cancellative elements, then we say that R is a transfer Krull domain (of finite type) if R^\bullet is a transfer Krull monoid (of finite type). Let $\theta: H \rightarrow B$ be a weak transfer homomorphism between atomic monoids. It is easy to show that for all $a \in H$ we have $L_H(a) = L_B(\theta(a))$ and hence $\mathcal{L}(H) = \mathcal{L}(B)$. Since monoids of zero-sum sequences are BF-monoids, the same is true for transfer Krull monoids.

Let H^* be a commutative Krull monoid (i.e., H^* is commutative, completely integrally closed, and v -noetherian). Then there is a weak transfer homomorphism $\beta: H^* \rightarrow \mathcal{B}(G_0)$ where G_0 is a subset of the class group of H^* . Since monoids of zero-sum sequences are commutative Krull monoids and since the composition of weak transfer homomorphisms is a weak transfer homomorphism again, a monoid is a transfer Krull monoid if and only if it allows a weak transfer homomorphism to a commutative Krull monoid. In particular, commutative Krull monoids are transfer Krull monoids. However, a transfer Krull monoid need neither be commutative nor v -noetherian nor completely integrally closed. To give a noncommutative example,

consider a bounded HNP (hereditary noetherian prime) ring R . If every stably free left R -ideal is free, then its multiplicative monoid of cancellative elements is a transfer Krull monoid ([34]). A class of commutative weakly Krull domains which are transfer Krull but not Krull will be given in Theorem 5.8. Extended lists of commutative Krull monoids and of transfer Krull monoids, which are not commutative Krull, are given in [17].

The next proposition summarizes some key results on the structure of sets of lengths of transfer Krull monoids.

Proposition 3.2.

1. *Every transfer Krull monoid of finite type satisfies the Structure Theorem for Sets of Lengths.*
2. *For every $M \in \mathbb{N}_0$ and every finite nonempty set $\Delta \subset \mathbb{N}$, there is a finite abelian group G such that the following holds: for every AAMP L with difference $d \in \Delta$ and bound M there is some $y_L \in \mathbb{N}$ such that*

$$y + L \in \mathcal{L}(G) \quad \text{for all } y \geq y_L.$$

3. *If G is an infinite abelian group, then*

$$\mathcal{L}(G) = \{L \subset \mathbb{N}_{\geq 2} \mid L \text{ is finite and nonempty}\} \cup \{\{0\}, \{1\}\}.$$

Proof. 1. Let H be a transfer Krull monoid and $\theta: H \rightarrow \mathcal{B}(G_0)$ be a weak transfer homomorphism where G_0 is a finite subset of an abelian group. Then $\mathcal{L}(H) = \mathcal{L}(G_0)$, and $\mathcal{B}(G_0)$ satisfies the Structure Theorem by [20, Theorem 4.4.11].

For 2. we refer to [33], and for 3. see [31] and [20, Section 7.4]. \square

The inequalities in (1) and the subsequent remarks show that a finite abelian group G has Davenport constant $D(G) \leq 4$ if and only if G is cyclic of order $|G| \leq 4$ or if it is isomorphic to $C_2 \oplus C_2$ or to C_2^3 . For these groups an explicit description of their systems of sets of lengths has been given, and we gather this in the next proposition (in Section 4 we will determine the systems $\mathcal{L}(G)$ for all groups G with $D(G) = 5$).

Proposition 3.3.

1. *If G is an abelian group, then $\mathcal{L}(G) = \{y + L \mid y \in \mathbb{N}_0, L \in \mathcal{L}(G^\bullet)\} \supset \{\{y\} \mid y \in \mathbb{N}_0\}$, and equality holds if and only if $|G| \leq 2$.*
2. $\mathcal{L}(C_3) = \mathcal{L}(C_2 \oplus C_2) = \{y + 2k + [0, k] \mid y, k \in \mathbb{N}_0\}.$
3. $\mathcal{L}(C_4) = \{y + k + 1 + [0, k] \mid y, k \in \mathbb{N}_0\} \cup \{y + 2k + 2 \cdot [0, k] \mid y, k \in \mathbb{N}_0\}.$
4. $\mathcal{L}(C_2^3) = \{y + (k+1) + [0, k] \mid y \in \mathbb{N}_0, k \in [0, 2]\}$
 $\cup \{y + k + [0, k] \mid y \in \mathbb{N}_0, k \geq 3\} \cup \{y + 2k + 2 \cdot [0, k] \mid y, k \in \mathbb{N}_0\}.$

Proof. See [20, Proposition 7.3.1 and Theorem 7.3.2]. \square

Let G and G' be abelian groups. Then their monoids of zero-sum sequences $\mathcal{B}(G)$ and $\mathcal{B}(G')$ are isomorphic if and only if the groups G and G' are isomorphic ([20, Corollary 2.5.7]). The standing conjecture states that the systems of sets of lengths $\mathcal{L}(G)$ and $\mathcal{L}(G')$ of finite groups coincide if and only if G and G' are isomorphic (apart from the trivial cases listed in Proposition 3.3). Here is the precise formulation of the conjecture (it was first stated in [17]).

Conjecture 3.4. *Let G be a finite abelian group with $D(G) \geq 4$. If G' is an abelian group with $\mathcal{L}(G) = \mathcal{L}(G')$, then G and G' are isomorphic.*

The conjecture holds true for groups G having rank $r(G) \leq 2$, for groups of the form $G = C_n^r$ (if r is small with respect to n), and others ([24, 27, 36]). But it is far open in general, and the goal of this section is to develop new viewpoints of looking at this conjecture.

Let G be a finite abelian group with $D(G) \geq 4$. If G' is a finite abelian group with $\mathcal{L}(G) = \mathcal{L}(G')$, then (2) shows that

$$\begin{aligned} D(G) &= \rho_2(G) = \sup\{\sup L \mid 2 \in L \in \mathcal{L}(G)\} \\ &= \sup\{\sup L \mid 2 \in L \in \mathcal{L}(G')\} = \rho_2(G') = D(G'). \end{aligned}$$

We see from the inequalities in (1) that there are (up to isomorphism) only finitely many finite abelian groups G' with given Davenport constant, and hence there are only finitely many finite abelian groups G' with $\mathcal{L}(G) = \mathcal{L}(G')$. Thus Conjecture 3.4 is equivalent to the statement that for each $m \geq 4$ and for each two non-isomorphic finite abelian groups G and G' having Davenport constant $D(G) = D(G') = m$ the systems $\mathcal{L}(G)$ and $\mathcal{L}(G')$ are distinct. Therefore we have to study the set

$$\Omega_m = \{\mathcal{L}(G) \mid G \text{ is a finite abelian group with } D(G) = m\}$$

of all systems of sets of lengths stemming from groups having Davenport constant equal to m . If a group G' is a proper subgroup of G , then $D(G') < D(G)$ ([20, Proposition 5.1.11]) and hence $\mathcal{L}(G') \subsetneq \mathcal{L}(G)$. Thus if $D(G) = D(G')$ for some group G' , then none of the groups is isomorphic to a proper subgroup of the other one. Conversely, if G' is a finite abelian group with $\mathcal{L}(G') \subset \mathcal{L}(G)$, then $D(G') = \rho_2(G') \leq \rho_2(G) = D(G)$. However, it may happen that $\mathcal{L}(G') \subsetneq \mathcal{L}(G)$ but $D(G') = D(G)$. Indeed, Proposition 3.3 shows that $\mathcal{L}(C_4) \subsetneq \mathcal{L}(C_2^3)$, and we will observe this phenomenon again in Section 4.

Theorem 3.5. *For $m \in \mathbb{N}$, let*

$$\Omega_m = \{\mathcal{L}(G) \mid G \text{ is a finite abelian group with } D(G) = m\}.$$

Then $\mathcal{L}(C_2^{m-1})$ is a maximal element and $\mathcal{L}(C_m)$ is a minimal element in Ω_m (with respect to set-theoretical inclusion). Furthermore, if G is an abelian group with $D(G) = m$ and $\mathcal{L}(G) \subset \mathcal{L}(C_2^{m-1})$, then $G \cong C_m$ or $G \cong C_2^{m-1}$.

Proof. If $m \in [1, 2]$, then $|\Omega_m| = 1$ and hence all assertions hold. Since C_3 and $C_2 \oplus C_2$ are the only groups (up to isomorphism) with Davenport constant three, and since $\mathcal{L}(C_3) = \mathcal{L}(C_2^2)$ by Proposition 3.3, the assertions follow. We suppose that $m \geq 4$ and proceed in two steps.

1. To show that $\mathcal{L}(C_2^{m-1})$ is maximal, we study, for a finite abelian group G , the set $\Delta_1(G)$. We define

$$\Delta^*(G) = \{\min \Delta(G_0) \mid G_0 \subset G \text{ with } \Delta(G_0) \neq \emptyset\},$$

and recall that (see [20, Corollary 4.3.16])

$$\Delta^*(G) \subset \Delta_1(G) \subset \{d_1 \in \Delta(G) \mid d_1 \text{ divides some } d \in \Delta^*(G)\}.$$

Thus $\max \Delta_1(G) = \max \Delta^*(G)$, and [29, Theorem 1.1] implies that $\max \Delta^*(G) = \max \{\exp(G) - 2, r(G) - 1\}$. Assume to the contrary that there is a finite abelian group G with $D(G) = m \geq 4$ that is not an elementary 2-group such that $\mathcal{L}(C_2^{m-1}) \subset \mathcal{L}(G)$. Then

$$\begin{aligned} m - 2 &= \max \Delta^*(C_2^{m-1}) = \max \Delta_1(C_2^{m-1}) \leq \max \Delta_1(G) \\ &= \max \Delta^*(G) = \max \{\exp(G) - 2, r(G) - 1\}. \end{aligned}$$

If $r(G) \geq m - 1$, then $D(G) = m$ implies that $G \cong C_2^{m-1}$, a contradiction. Thus $\exp(G) \geq m$, and since $D(G) = m$ we infer that $G \cong C_m$. If $m = 4$, then Proposition 3.3.4 shows that $\mathcal{L}(C_2^3) \not\subset \mathcal{L}(C_4)$, a contradiction. Suppose that $m \geq 5$. Then $\Delta^*(C_2^{m-1}) = \Delta_1(C_2^{m-1}) = \Delta(C_2^{m-1}) = [1, m-2]$ by [20, Corollary 6.8.3]. For cyclic groups we have $\max \Delta^*(C_m) = m - 2$ and $\max(\Delta^*(C_m) \setminus \{m-2\}) = \lfloor m/2 \rfloor - 1$ by [20, Theorem 6.8.12]. Therefore $\mathcal{L}(C_2^{m-1}) \subset \mathcal{L}(C_m)$ implies that

$$[1, m-2] = \Delta_1(C_2^{m-1}) \subset \Delta_1(C_m),$$

a contradiction to $m-3 \notin \Delta_1(C_m)$.

2. We recall some facts. Let G be a group with $D(G) = m$. If $U \in \mathcal{A}(G)$ with $|U| = D(G)$, then $\{2, D(G)\} \subset L(U(-U))$. Cyclic groups and elementary 2-groups are the only groups G with the following property: if $L \in \mathcal{L}(G)$ with $\{2, D(G)\} \subset L$, then $L = \{2, D(G)\}$ ([20, Theorem 6.6.3]).

Now assume to the contrary that there is a finite abelian group G with $D(G) = m$ such that $\mathcal{L}(G) \subset \mathcal{L}(C_m)$. Let $L \in \mathcal{L}(G)$ with $\{2, D(G)\} \subset L$. Then $L \in \mathcal{L}(C_m)$ whence $L = \{2, D(G)\}$ which implies that G is cyclic or an elementary 2-group. By 1., G is not an elementary 2-group whence G is cyclic which implies $G \cong C_m$ and hence $\mathcal{L}(G) = \mathcal{L}(C_m)$.

The furthermore assertion on groups G with $D(G) = m$ and $\mathcal{L}(G) \subset \mathcal{L}(C_2^{m-1})$ follows as above by considering sets of lengths L with $\{2, D(G)\} \subset L$. \square

In Section 4 we will see that $\mathcal{L}(C_2^{m-1})$ need not be the largest element in Ω_m , and that indeed $\mathcal{L}(C_m) \subset \mathcal{L}(C_2^{m-1})$ for $m \in [2, 5]$, where the inclusion is strict for

$m \geq 4$. On the other hand, it is shown in [28] that $\mathcal{L}(C_m) \not\subset \mathcal{L}(C_2^{m-1})$ for infinitely many $m \in \mathbb{N}$.

Theorem 3.6. *We have*

$$\bigcap \mathcal{L}(G) = \{y + 2k + [0, k] \mid y, k \in \mathbb{N}_0\},$$

where the intersection is taken over all finite abelian groups G with $|G| \geq 3$.

Proof. By Proposition 3.3.2, the intersection on the left hand side is contained in the set on the right hand side. Let G be a finite abelian group with $|G| \geq 3$. If $L \in \mathcal{L}(G)$, then $y + L \in \mathcal{L}(G)$. Thus it is sufficient to show that $[2k, 3k] \in \mathcal{L}(G)$ for every $k \in \mathbb{N}$. If G contains two independent elements of order 2 or an element of order 4, then the claim follows by Proposition 3.3. Thus, it remains to consider the case when G contains an element g with $\text{ord}(g) = p$ for some odd prime $p \in \mathbb{N}$. Let $k \in \mathbb{N}$ and $B_k = ((2g)^p g^p)^k$. We assert that $\mathcal{L}(B_k) = [2k, 3k]$.

We set $U_1 = g^p$, $U_2 = (2g)^p$, $V_1 = (2g)^{(p-1)/2}g$, and $V_2 = (2g)g^{p-2}$. Since $U_1 U_2 = V_1^2 V_2$ and

$$B_k = (U_1 U_2)^k = (U_1 U_2)^{k-v} (V_1^2 V_2)^v \quad \text{for all } v \in [0, k],$$

it follows that $[2k, 3k] \subset \mathcal{L}(B_k)$.

In order to show there are no other factorization lengths, we recall the concept of the g -norm of sequences. If $S = (n_1 g) \cdot \dots \cdot (n_\ell g) \in \mathcal{B}(\langle g \rangle)$, where $\ell \in \mathbb{N}_0$ and $n_1, \dots, n_\ell \in [1, \text{ord}(g)]$, then

$$\|S\|_g = \frac{n_1 + \dots + n_\ell}{\text{ord}(g)} \in \mathbb{N}$$

is the g -norm of S . Clearly, if $S = S_1 \cdot \dots \cdot S_m$ with $S_1, \dots, S_m \in \mathcal{A}(G)$, then $\|S\|_g = \|S_1\|_g + \dots + \|S_m\|_g$.

Note that $U_2 = (2g)^p$ is the only atom in $\mathcal{A}(\{g, 2g\})$ with g -norm 2, and all other atoms in $\mathcal{A}(\{g, 2g\})$ have g -norm 1. Let $B_k = W_1 \cdot \dots \cdot W_\ell$ be a factorization of B_k , and let ℓ' be the number of $i \in [1, \ell]$ such that $W_i = (2g)^p$. We have $\|B_k\|_g = 3k$ and thus $3k = 2\ell' + (\ell - \ell') = \ell' + \ell$. Since $\ell' \in [0, k]$, it follows that $\ell = 3k - \ell' \in [2k, 3k]$. \square

Theorem 3.7. *Let $L \subset \mathbb{N}_{\geq 2}$ be a finite nonempty subset. Then there are only finitely many pairwise non-isomorphic finite abelian groups G such that $L \notin \mathcal{L}(G)$.*

Proof. We start with the following two assertions.

- A1.** There is an integer $n_L \in \mathbb{N}$ such that $L \in \mathcal{L}(C_n)$ for every $n \geq n_L$.
- A2.** For every $p \in \mathbb{P}$ there is an integer $r_{p,L} \in \mathbb{N}$ such that $L \in \mathcal{L}(C_p^r)$ for every $r \geq r_{p,L}$.

Proof of A1. By Proposition 3.2.3, there is some $B = \prod_{i=1}^k m_i \prod_{j=1}^{\ell} (-n_j) \in \mathcal{B}(\mathbb{Z})$ such that $L(B) = L$, where $k, \ell, m_1, \dots, m_k \in \mathbb{N}$ and $n_1, \dots, n_{\ell} \in \mathbb{N}_0$. We set $n_L = n_1 + \dots + n_{\ell}$ and choose some $n \in \mathbb{N}$ with $n \geq n_L$. If $S \in \mathcal{F}(\mathbb{Z})$ with $S|B$ and $f: \mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$ denotes the canonical epimorphism, then S has sum zero if and only if $f(S)$ has sum zero. This implies that $L_{\mathcal{B}(\mathbb{Z}/n\mathbb{Z})}(f(B)) = L_{\mathcal{B}(\mathbb{Z})}(B) = L$. \square

[Proof of A1]

Proof of A2. Let $p \in \mathbb{P}$ be a prime and let G_p be an infinite dimensional \mathbb{F}_p -vector space. By Proposition 3.2.3, there is some $B_p \in \mathcal{B}(G_p)$ such that $L(B_p) = L$. If $r_{p,L}$ is the rank of $\langle \text{supp}(B_p) \rangle \subset G_p$, then

$$L = L(B_p) \in \mathcal{L}(\langle \text{supp}(B_p) \rangle) \subset \mathcal{L}(C_p^r) \quad \text{for } r \geq r_{p,L}. \quad \square[\text{Proof of A2}]$$

Now let G be a finite abelian group such that $L \notin \mathcal{L}(G)$. Then **A1** implies that $\exp(G) < n_L$, and **A2** implies that $r_p(G) < r_{p,L}$ for all primes p with $p \mid \exp(G)$. Thus the assertion follows. \square

4 Sets of lengths of transfer Krull monoids over small groups

Since the very beginning of factorization theory, invariants controlling the structure of sets of lengths (such as elasticities and sets of distances) have been in the center of interest. Nevertheless, (apart from a couple of trivial cases) the full system of sets of lengths has been written down explicitly only for the following classes of monoids:

- Numerical monoids generated by arithmetical progressions: see [1].
- Self-idealizations of principal ideal domains: see [10, Corollary 4.16], [4, Remark 4.6].
- The ring of integer-valued polynomials over \mathbb{Z} : see [15].
- The systems $\mathcal{L}(G)$ for infinite abelian groups G and for abelian groups G with $D(G) \leq 4$: see Propositions 3.2 and 3.3.

The goal of this section is to determine $\mathcal{L}(G)$ for abelian groups G having Davenport constant $D(G) = 5$. By inequality (1) and the subsequent remarks, a finite abelian group G has Davenport constant five if and only if it is isomorphic to one of the following groups:

$$C_3 \oplus C_3, \quad C_5, \quad C_2 \oplus C_4, \quad C_2^4.$$

Their systems of sets of lengths are given in Theorems 4.1, 4.3, 4.5, and 4.8. We start with a brief analysis of these explicit descriptions (note that they will be needed again in Section 5; confer the proof of Theorem 5.7).

By Theorem 3.5, we know that $\mathcal{L}(C_2^4)$ is maximal in $\Omega_5 = \{\mathcal{L}(C_5), \mathcal{L}(C_2 \oplus C_4), \mathcal{L}(C_3 \oplus C_3), \mathcal{L}(C_2^4)\}$. Theorems 4.1, 4.3, 4.5, and 4.8 unveil that $\mathcal{L}(C_3 \oplus C_3)$,

$\mathcal{L}(C_2 \oplus C_4)$, and $\mathcal{L}(C_2^4)$ are maximal in Ω_5 , and that $\mathcal{L}(C_5)$ is contained in $\mathcal{L}(C_2^4)$, but it is neither contained in $\mathcal{L}(C_3 \oplus C_3)$ nor in $\mathcal{L}(C_2 \oplus C_4)$. Furthermore, Theorems 3.5, 4.3, and 4.8 show that $\mathcal{L}(C_m) \subset \mathcal{L}(C_2^{m-1})$ for $m \in [2, 5]$. It is well-known that, for all $m \geq 4$, $\mathcal{L}(C_m) \neq \mathcal{L}(C_2^{m-1})$ ([16, Corollary 5.3.3]), and the standing conjecture is that $\mathcal{L}(C_m) \not\subset \mathcal{L}(C_2^{m-1})$ holds true for almost all $m \in \mathbb{N}_{\geq 2}$ (see [28]).

The group $C_3 \oplus C_3$ has been handled in [24, Theorem 4.2].

Theorem 4.1. $\mathcal{L}(C_3^2) = \{y + [2k, 5k] \mid y, k \in \mathbb{N}_0\} \cup \{y + [2k+1, 5k+2] \mid y \in \mathbb{N}_0, k \in \mathbb{N}\}$.

Remark. An equivalent way to describe $\mathcal{L}(C_3^2)$ is $\{y + \lceil \frac{2k}{3} \rceil + [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N}_{\geq 2}\} \cup \{\{y\}, y+2+[0, 1] \mid y \in \mathbb{N}_0\}$.

The fact that all sets of lengths are intervals is a consequence of the fact $\Delta(C_3^2) = \{1\}$. Of course, each set of lengths L has to fulfill $\rho(L) \leq 5/2 = \rho(C_3^2)$. We observe that the description shows that this is the only condition, provided $\min L \geq 2$. The following lemma is frequently helpful in the remainder of this section.

Lemma 4.2. *Let G be a finite abelian group, and let $A \in \mathcal{B}(G)$.*

1. *If $\text{supp}(A) \cup \{0\}$ is a group, then $\mathcal{L}(A)$ is an interval.*
2. *If A_1 is an atom dividing A with $|A_1| = 2$, then $\max \mathcal{L}(A) = 1 + \max \mathcal{L}(AA_1^{-1})$.*
3. *If A is a product of atoms of length 2 and if every atom A_1 dividing A has length $|A_1| = 2$ or $|A_1| = 4$, then $\max \mathcal{L}(A) - 1 \notin \mathcal{L}(A)$.*

Proof. 1. See [20, Theorem 7.6.8].

2. Let $\ell = \max \mathcal{L}(A)$ and $A = U_1 \cdot \dots \cdot U_\ell$, where $U_1, \dots, U_\ell \in \mathcal{A}(G)$. Let $A_1 = g_1 g_2$, where $g_1, g_2 \in G$. If there exists $i \in [1, \ell]$ such that $A_1 = U_i$, then $\max \mathcal{L}(A) = 1 + \max \mathcal{L}(AA_1^{-1})$. Otherwise there exist distinct $i, j \in [1, \ell]$ such that $g_1 \mid U_i$ and $g_2 \mid U_j$. Thus A_1 divides $U_i U_j$ and hence $1 + \max \mathcal{L}(AA_1^{-1}) \geq \ell$ which implies that $\max \mathcal{L}(A) = 1 + \max \mathcal{L}(AA_1^{-1})$ by the maximality of ℓ .

3. If $\max \mathcal{L}(A) - 1 \in \mathcal{L}(A)$, then $A = V_1 \cdot \dots \cdot V_{\max \mathcal{L}(A)-1}$ with $|V_1| = 4$ and $|V_2| = \dots = |V_{\max \mathcal{L}(A)-1}| = 2$. Thus V_1 can only be a product two atoms of length 2, a contradiction. \square

We now consider the groups C_5 , $C_2 \oplus C_4$, and C_2^4 , each one in its own subsection. In the proofs of the forthcoming theorems we will use Proposition 3.3 and Theorem 3.6 without further mention.

4.1 The system of sets of lengths of C_5

The goal of this subsection is to prove the following result.

Theorem 4.3. $\mathcal{L}(C_5) = \mathcal{L}_1 \cup \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5 \cup \mathcal{L}_6$, where

$$\begin{aligned}\mathcal{L}_1 &= \{\{y\} \mid y \in \mathbb{N}_0\}, \\ \mathcal{L}_2 &= \{y+2 + \{0, 2\} \mid y \in \mathbb{N}_0\}, \\ \mathcal{L}_3 &= \{y+3 + \{0, 1, 3\} \mid y \in \mathbb{N}_0\}, \\ \mathcal{L}_4 &= \{y+2k+3 \cdot [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N}\}, \\ \mathcal{L}_5 &= \{y+2 \left\lceil \frac{k}{3} \right\rceil + [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N} \setminus \{3\}\} \cup \{y+[3, 6] \mid y \in \mathbb{N}_0\}, \text{ and} \\ \mathcal{L}_6 &= \{y+2k+3 + \{0, 2, 3\} + 3 \cdot [0, k] \mid y, k \in \mathbb{N}_0\}.\end{aligned}$$

We observe that all sets of lengths with many elements are arithmetic multi-progressions with difference 1 or 3. Yet, there are none with difference 2. This is because $\Delta^*(C_5) = \{1, 3\}$. Moreover, we point out that the condition for an interval to be a set of lengths is different from that of the other groups with Davenport constant 5. This is related to the fact that $\rho_{2k+1}(C_5) = 5k+1$, while $\rho_{2k+1}(G) = 5k+2$ for the other groups with Davenport constant 5. Before we start the actual proof, we collect some results on sets of lengths over C_5 .

Lemma 4.4. Let G be cyclic of order five, and let $A \in \mathcal{B}(G)$.

1. If $g \in G^\bullet$ and $k \in \mathbb{N}_0$, then

$$L(g^{5(k+1)}(-g)^{5(k+1)}(2g)g^3) = 2k+3 + \{0, 2, 3\} + 3 \cdot [0, k].$$

2. If $2 \in \Delta(L(A)) \subset [1, 2]$, then $L(A) \in \{\{y, y+2\} \mid y \geq 2\} \cup \{\{y, y+1, y+3\} \mid y \geq 3\}$ or $L(A) = 3 + \{0, 2, 3\} + L(A')$ where $A' \in \mathcal{B}(G)$ and $L(A')$ is an arithmetical progression of difference 3.
3. $\Delta(G) = [1, 3]$, and if $3 \in \Delta(L(A))$, then $\Delta(L(A)) = \{3\}$.
4. $\rho_{2k+1}(G) = 5k+1$ for all $k \in \mathbb{N}$.

Proof. 1. and 2. follow from the proof of [24, Lemma 4.5].

3. See [20, Theorems 6.7.1 and 6.4.7] and [12, Theorem 3.3].

4. See [16, Theorem 5.3.1]. \square

Proof (Theorem 4.3). Let G be cyclic of order five and let $g \in G^\bullet$. We first show that all the specified sets occur as sets of lengths, and then we show that no other sets occur.

Step 1. We prove that for every $L \in \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5 \cup \mathcal{L}_6$, there exists an $A \in \mathcal{B}(G)$ such that $L = L(A)$. We distinguish five cases.

If $L = \{y, y+2\} \in \mathcal{L}_2$ with $y \geq 2$, then we set $A = 0^{y-2}g^5(-g)^3(-2g)$ and obtain that $L(A) = y-2 + \{2, 4\} = L$.

If $L = \{y, y+1, y+3\} \in \mathcal{L}_3$ with $y \geq 3$, then we set $A = 0^{y-3}g^5(-g)^5g^2(-2g)$ and obtain that $L(A) = y-3 + \{3, 4, 6\} = \{y, y+1, y+3\} = L$.

If $L = y+2k+3 \cdot [0, k] \in \mathcal{L}_4$ with $k \in \mathbb{N}$ and $y \in \mathbb{N}_0$, then we set $A = g^{5k}(-g)^{5k}0^y \in \mathcal{B}(G)$ and hence $L(A) = y+2k+3 \cdot [0, k] = L$.

If $L = y + 2k + 3 + \{0, 2, 3\} + 3 \cdot [0, k] \in \mathcal{L}_6$ with $k \in \mathbb{N}_0$ and $y \in \mathbb{N}_0$, then we set $A = 0^y g^{5(k+1)} (-g)^{5(k+1)} (2g) g^3$ and hence $L(A) = y + 2k + 3 + \{0, 2, 3\} + 3 \cdot [0, k] = L$ by Lemma 4.4.1.

Now we suppose that $L \in \mathcal{L}_5$, and we distinguish two subcases. First, if $L = y + [3, 6]$ with $y \in \mathbb{N}_0$, then we set $A = 0^y (2g(-2g)) g^5 (-g)^5$ and hence $L(A) = y + [3, 6] = L$. Second, we assume that $L = y + 2 \lceil \frac{k}{3} \rceil + [0, k]$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N} \setminus \{3\}$.

If $k \in \mathbb{N}$ with $k \equiv 0 \pmod{3}$, then $k \geq 6$ and by Lemma 4.2.1 we obtain that

$$L(0^y (2g)^5 (-2g)^5 g^{5t} (-g)^{5t}) = y + [2t + 2, 5t + 5] = y + 2 \lceil \frac{k}{3} \rceil + [0, k] = L,$$

where $k = 3t + 3$.

If $k \in \mathbb{N}$ with $k \equiv 1 \pmod{3}$, then by Lemma 4.2.1 we obtain that

$$L(0^y (2g(-g)^2) (g^2(-2g)) g^{5t} (-g)^{5t}) = y + [2t + 2, 5t + 3] = y + 2 \lceil \frac{k}{3} \rceil + [0, k] = L,$$

where $k = 3t + 1$.

If $k \in \mathbb{N}$ with $k \equiv 2 \pmod{3}$, then by Lemma 4.2.1 we obtain that

$$L(0^y (g^3(2g))((-g)^3(-2g)) g^{5t} (-g)^{5t}) = y + [2t + 2, 5t + 4] = y + 2 \lceil \frac{k}{3} \rceil + [0, k] = L,$$

where $k = 3t + 2$.

Step 2. We prove that for every $A \in \mathcal{B}(G^\bullet)$, $L(A) \in \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5 \cup \mathcal{L}_6$.

Let $A \in \mathcal{B}(G^\bullet)$. We may suppose that $\Delta(L(A)) \neq \emptyset$. By Lemma 4.4.3 we distinguish three cases according to the form of the set of distances $\Delta(L(A))$.

CASE 1: $\Delta(L(A)) = \{1\}$.

Then $L(A)$ is an interval and hence we assume that $L(A) = [y, y+k] = y + [0, k]$ where $y \geq 2$ and $k \geq 1$. If $k = 3$ and $y = 2$, then $L(A) = [2, 5]$ and hence $L(A) = L(g^5(-g)^5) = \{2, 5\}$, a contradiction. Thus $k = 3$ implies that $y \geq 3$ and hence $L(A) \in \mathcal{L}_5$. If $k \leq 2$, then we obviously have that $L(A) \in \mathcal{L}_5$. Suppose that $k \geq 4$. If $y = 2t$ with $t \geq 2$, then $y+k \leq 5t$ and hence $y = 2t \geq 2 \lceil \frac{k}{3} \rceil$ which implies that $L(A) \in \mathcal{L}_5$. If $y = 2t+1$ with $t \geq 1$, then $y+k \leq 5t+1$ and hence $y = 2t+1 \geq 1 + 2 \lceil \frac{k}{3} \rceil$ which implies that $L(A) \in \mathcal{L}_5$.

CASE 2: $\Delta(L(A)) = \{3\}$.

Then $L(A) = y + 3 \cdot [0, k]$ where $y \geq 2$ and $k \geq 1$. If $y = 2t \geq 2$, then $y+3k \leq 5t$ and hence $y = 2t \geq 2k$ which implies that $L(A) \in \mathcal{L}_4$. If $y = 2t+1 \geq 3$, then $y+3k \leq 5t+1$ and hence $y = 2t+1 \geq 1+2k$ which implies that $L(A) \in \mathcal{L}_4$.

CASE 3: $2 \in \Delta(L(A)) \subset [1, 2]$.

By Lemma 4.4.2, we infer that either $L(A) \in \mathcal{L}_2 \cup \mathcal{L}_3$ or that $L(A) = 3 + \{0, 2, 3\} + L(A')$, where $A' \in \mathcal{B}(G)$ and $L(A')$ is an arithmetical progression of difference 3. In the latter case we obtain that $L(A') = y + 2k + 3 \cdot [0, k]$, with $y \in \mathbb{N}_0$ and $k \in \mathbb{N}_0$, and hence $L(A) = y + 2k + 3 + \{0, 2, 3\} + 3 \cdot [0, k] \in \mathcal{L}_6$. \square

4.2 The system of sets of lengths of $C_2 \oplus C_4$

We establish the following result, giving a complete description of the system of sets of lengths of $C_2 \oplus C_4$.

Theorem 4.5. $\mathcal{L}(C_2 \oplus C_4) = \mathcal{L}_1 \cup \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5$, where

$$\begin{aligned}\mathcal{L}_1 &= \{\{y\} \mid y \in \mathbb{N}_0\}, \\ \mathcal{L}_2 &= \left\{y + 2 \left\lceil \frac{k}{3} \right\rceil + [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N} \setminus \{3\}\right\} \cup \\ &\quad \{y + [3, 6] \mid y \in \mathbb{N}_0\} \cup \{[2t + 1, 5t + 2] \mid t \in \mathbb{N}\} \\ &= \left\{y + \left\lceil \frac{2k}{3} \right\rceil + [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N} \setminus \{1, 3\}\right\} \cup \\ &\quad \{y + 3 + [0, 3], y + 2 + [0, 1] \mid y \in \mathbb{N}_0\}, \\ \mathcal{L}_3 &= \{y + 2k + 2 \cdot [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N}\}, \\ \mathcal{L}_4 &= \{y + k + 1 + (\{0\} \cup [2, k + 2]) \mid y \in \mathbb{N}_0, k \in \mathbb{N} \text{ odd}\}, \text{ and} \\ \mathcal{L}_5 &= \{y + k + 2 + ([0, k] \cup \{k + 2\}) \mid y \in \mathbb{N}_0, k \in \mathbb{N}\}.\end{aligned}$$

We note that all sets of lengths are arithmetical progressions with difference 2 or almost arithmetical progressions with difference 1 and bound 2. This is related to the fact that $\Delta(C_2 \oplus C_4) = \Delta^*(C_2 \oplus C_4) = \{1, 2\}$. We start with a lemma determining all minimal zero-sum sequences over $C_2 \oplus C_4$.

Lemma 4.6. Let (e, g) be a basis of $G = C_2 \oplus C_4$ with $\text{ord}(e) = 2$ and $\text{ord}(g) = 4$. Then the minimal zero-sum sequences over G^\bullet are given by the following list.

1. The minimal zero sum sequences of length 2 are :

$$\begin{aligned}S_2^1 &= \{e^2, (e + 2g)^2\}, \\ S_2^2 &= \{(2g)^2\}, \\ S_2^3 &= \{g(-g), (e + g)(e - g)\}\end{aligned}$$

2. The minimal zero sum sequences of length 3 are :

$$\begin{aligned}S_3^1 &= \{e(2g)(e + 2g)\}, \\ S_3^2 &= \{g^2(2g), (-g)^2(2g), (e + g)^2(2g), (e - g)^2(2g)\}, \\ S_3^3 &= \{eg(e - g), e(-g)(e + g), (e + 2g)g(e + g), (e + 2g)(-g)(e - g)\}.\end{aligned}$$

3. The minimal zero sum sequences of length 4 are :

$$\begin{aligned}
S_4^1 &= \{g^4, (-g)^4, (e+g)^4, (e-g)^4\}, \\
S_4^2 &= \{g^2(e+g)^2, (-g)^2(e-g)^2, g^2(e-g)^2, (-g)^2(e+g)^2\}, \\
S_4^3 &= \{eg^2(e+2g), e(e+g)^2(e+2g), e(-g)^2(e+2g), e(e-g)^2(e+2g)\}, \\
S_4^4 &= \{eg(2g)(e+g), e(-g)(2g)(e-g), (e+2g)g(2g)(e-g), (e+2g)(-g)(2g)(e+g)\}.
\end{aligned}$$

4. The minimal zero sum sequences of length 5 are :

$$\begin{aligned}
S_5 = \{ &eg^3(e+g), e(-g)^3(e-g), e(e+g)^3g, e(e-g)^3(-g) \\
&(e+2g)g^3(e-g), (e+2g)(-g)^3(e+g), \\
&(e+2g)(e+g)^3(-g), (e+2g)(e-g)^3g \},
\end{aligned}$$

Moreover, for each two atoms W_1, W_2 in any one of the above sets, there exists a group isomorphism $\phi: G \rightarrow G$ such that $\phi(W_1) = W_2$.

Proof. We give a sketch of the proof.

Since a minimal zero-sum sequence of length two is of the form $h(-h)$ for some non-zero element $h \in G$, the list given in 1. follows.

A minimal zero-sum sequence of length three contains either two elements of order four or no element of order four. If there are two elements of order four, we can have one element of order four with multiplicity two (see S_3^2) or two distinct elements of order four that are not the inverse of each other (see S_3^3). If there is no element of order four, the sequence consists of three distinct elements of order two (see S_3^1).

A minimal zero-sum sequence of length four contains either four elements of order four or two elements of order four. If there are two elements of order four, the sequence can contain one element with multiplicity two (see S_4^3) or any two distinct elements that are not each other's inverse with multiplicity one (see S_4^4). If there are four elements of order four, the sequence can contain one element with multiplicity four (see S_4^1) or two elements with multiplicity two (see S_4^2).

Since every minimal zero-sum sequence of length five contains an element with multiplicity three, the list given in 4. follows (for details see [20, Theorem 6.6.5]).

The existence of the required isomorphism follows immediately from the given description of the sequences. \square

The next lemma collects some basic results on $\mathcal{L}(C_2 \oplus C_4)$ that will be essential for the proof of Theorem 4.5.

Lemma 4.7. *Let $G = C_2 \oplus C_4$, and let $A \in \mathcal{B}(G)$.*

1. $\Delta(G) = [1, 2]$, and if $\{2, 5\} \subset \mathsf{L}(A)$, then $\mathsf{L}(A) = \{2, 4, 5\}$.
2. $\rho_{2k+1}(G) = 5k + 2$ for all $k \in \mathbb{N}$.

3. If (e, g) is a basis of G with $\text{ord}(e) = 2$ and $\text{ord}(g) = 4$, then $\{0, g, 2g, e+g, e+2g\}$ and $\{0, g, 2g, e, e-g\}$ are half-factorial sets. Furthermore, if $\text{supp}(A) \subset \{e, g, 2g, e+g, e+2g\}$ and $v_e(A) = 1$, then $|\mathcal{L}(A)| = 1$.

Proof. 1. The first assertion follows from [20, Theorem 6.7.1 and Corollary 6.4.8]. Let $A \in \mathcal{B}(G)$ with $\{2, 5\} \subset \mathcal{L}(A)$. Then there is an $U \in \mathcal{A}(G)$ of length $|U| = 5$ such that $A = (-U)U$. By Lemma 4.6 there is a basis (e, g) of G with $\text{ord}(e) = 2$ and $\text{ord}(g) = 4$ such that $U = eg^3(e+g)$. This implies that $\mathcal{L}(A) = \{2, 4, 5\}$.

2. See [19, Corollary 5.2].

3. See [20, Theorem 6.7.9.1] for the first statement. Suppose that $\text{supp}(A) \subset \{e, g, 2g, e+g, e+2g\}$ and $v_e(A) = 1$. Then for every atom W dividing A with $e \mid W$, we have that $k(W) = \frac{3}{2}$. Since $\text{supp}(AW^{-1})$ is half-factorial, we obtain that $\mathcal{L}(AW^{-1}) = \{k(A) - 3/2\}$ by [20, Proposition 6.7.3] which implies that $\mathcal{L}(A) = \{1 + k(A) - 3/2\} = \{k(A) - 1/2\}$. \square

Proof (Theorem 4.5). Let (e, g) be a basis of $G = C_2 \oplus C_4$ with $\text{ord}(e) = 2$ and $\text{ord}(g) = 4$. We start by collecting some basic constructions that will be useful. Then, we show that all the sets in the result actually are sets of lengths. Finally, we show there are no other sets of lengths.

Step 0. Some elementary constructions.

Let $U_1 = eg^3(e+g)$, $U_2 = (e+2g)(e+g)^3(-g)$, $U_3 = e(e-g)^3(-g)$, $U_4 = (-g)^2(e+g)^2$, and $U_5 = e(e+2g)g^2$. Then it is not hard to check that

$$\begin{aligned} \mathcal{L}(U_1(-U_1)) &= \mathcal{L}(U_2(-U_2)) = \{2, 4, 5\}, \\ \mathcal{L}(U_1U_3) &= [2, 4], & \mathcal{L}(U_1(-U_4)) &= [2, 3], \\ \mathcal{L}(U_1U_3U_4) &= [3, 7], & \mathcal{L}(U_1(-U_1)U_2(-U_2)) &= [4, 10], \\ \mathcal{L}(U_5^2(-g)^4) &= \{3, 4, 6\}, & \mathcal{L}(U_5(-U_5)g^4(-g)^4) &= \{4, 5, 6, 8\}, \\ \mathcal{L}(U_1(-U_1)(e+2g)^2) &= [3, 6]. \end{aligned} \tag{3}$$

Based on these results, we can obtain the sets of lengths of more complex zero-sum sequences. Let $k \in \mathbb{N}$.

Since $[2k+2, 4k+5] \supset \mathcal{L}(U_1(-U_1)g^{4k}(-g)^{4k}) \supset \mathcal{L}(U_1(-U_1)) + \mathcal{L}(g^{4k}(-g)^{4k}) = 2k+2 + (\{0\} \cup [2, 2k+3])$ and $2k+3 \notin \mathcal{L}(U_1(-U_1)g^{4k}(-g)^{4k})$, we obtain that

$$\mathcal{L}(U_1(-U_1)g^{4k}(-g)^{4k}) = 2k+2 + (\{0\} \cup [2, 2k+3]). \tag{4}$$

Since $[2(k+1), 5(k+1)] \supset \mathcal{L}(U_1(-U_1)U_2^k(-U_2)^k) \supset \mathcal{L}(U_1(-U_1)U_2(-U_2)) + \mathcal{L}(U_2^{k-1}(-U_2)^{k-1}) = [2(k+1), 5(k+1)]$, we obtain that

$$\mathcal{L}(U_1(-U_1)U_2^k(-U_2)^k) = [2(k+1), 5(k+1)]. \tag{5}$$

Since $[2(k+1), 5(k+1)-1] \supset \mathcal{L}(U_1U_3U_2^k(-U_2)^k) \supset \mathcal{L}(U_1U_3) + \mathcal{L}(U_2^k(-U_2)^k) = [2(k+1), 5(k+1)-1]$, we obtain that

$$\mathcal{L}(U_1U_3U_2^k(-U_2)^k) = [2(k+1), 5(k+1)-1]. \tag{6}$$

Since $[2(k+1), 5(k+1)-2] \supset \mathsf{L}(U_1(-U_4)U_2^k(-U_2)^k) \supset \mathsf{L}(U_1(-U_4)) + \mathsf{L}(U_2^k(-U_2)^k)$ and $\mathsf{L}(U_1(-U_4)) + \mathsf{L}(U_2^k(-U_2)^k) = [2(k+1), 5(k+1)-2]$, we obtain that

$$\mathsf{L}(U_1(-U_4)U_2^k(-U_2)^k) = [2(k+1), 5(k+1)-2]. \quad (7)$$

Since

$$[2k+1, 5k+2] \supset \mathsf{L}(U_1U_3U_4U_2^{k-1}(-U_2)^{k-1}) \supset \mathsf{L}(U_1U_3U_4) + \mathsf{L}(U_2^{k-1}(-U_2)^{k-1})$$

and $\mathsf{L}(U_1U_3U_4) + \mathsf{L}(U_2^{k-1}(-U_2)^{k-1}) = [2k+1, 5k+2]$, we obtain that

$$\mathsf{L}(U_1U_3U_4U_2^{k-1}(-U_2)^{k-1}) = [2k+1, 5k+2]. \quad (8)$$

Since

$$[2k+1, 4k+2] \supset \mathsf{L}(U_5^2(-g)^4g^{4k-4}(-g)^{4k-4}) \supset \mathsf{L}(U_5^2(-g)^4) + \mathsf{L}(g^{4k-4}(-g)^{4k-4}),$$

$$\mathsf{L}(U_5^2(-g)^4) + \mathsf{L}(g^{4k-4}(-g)^{4k-4}) = [2k+1, 4k] \cup \{4k+2\} \text{ and}$$

$$4k+1 \notin \mathsf{L}(U_5^2(-g)^4g^{4k-4}(-g)^{4k-4})$$

by Lemma 4.2.3, we obtain that

$$\mathsf{L}(U_5^2(-g)^4g^{4k-4}(-g)^{4k-4}) = [2k+1, 4k] \cup \{4k+2\}. \quad (9)$$

Suppose that $k \geq 2$. Since

$$[2k, 4k] \supset \mathsf{L}(U_5(-U_5)g^{4k-4}(-g)^{4k-4}) \supset \mathsf{L}(U_5(-U_5)g^4(-g)^4) + \mathsf{L}(g^{4k-8}(-g)^{4k-8}),$$

$$\mathsf{L}(U_5(-U_5)g^4(-g)^4) + \mathsf{L}(g^{4k-8}(-g)^{4k-8}) = [2k, 4k-2] \cup \{4k\}, \text{ and}$$

$$4k-1 \notin \mathsf{L}(U_5(-U_5)g^{4k-4}(-g)^{4k-4})$$

by Lemma 4.2.3, we obtain that

$$\mathsf{L}(U_5(-U_5)g^{4k-4}(-g)^{4k-4}) = [2k, 4k-2] \cup \{4k\}. \quad (10)$$

Step 1. We prove that for every $L \in \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5$ there exists an $A \in \mathcal{B}(G)$ such that $L = \mathsf{L}(A)$.

We distinguish four cases.

First we suppose that $L \in \mathcal{L}_2$, and we distinguish several subcases. If $L = y + [3, 6]$ with $y \in \mathbb{N}_0$, then we set $A = 0^y U_1(-U_1)(e+2g)^2$ and hence $\mathsf{L}(A) = y + [3, 6] = L$ by Equation (3). If $L = [2k+1, 5k+2]$ with $k \in \mathbb{N}$, then we set $A = U_1U_3U_4U_2^{k-1}(-U_2)^{k-1}$ and hence $\mathsf{L}(A) = L$ by Equation (8). Now we assume that $L = y + 2\lceil \frac{k}{3} \rceil + [0, k]$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N} \setminus \{3\}$.

If $k \equiv 0 \pmod{3}$, then $k \geq 6$ and by Equation (5) we infer that

$$L(0^y U_1(-U_1) U_2^t(-U_2)^t) = y + [2t+2, 5t+5] = y + 2\lceil \frac{k}{3} \rceil + [0, k] = L, \text{ where } k = 3t+3.$$

If $k \equiv 1 \pmod{3}$, then by Equation (7) we infer that

$$L(0^y U_1(-U_4) U_2^t(-U_2)^t) = y + [2t+2, 5t+3] = y + 2\lceil \frac{k}{3} \rceil + [0, k] = L, \text{ where } k = 3t+1.$$

If $k \equiv 2 \pmod{3}$, then by Equation (6) we infer that

$$L(0^y U_1 U_3 U_2^t(-U_2)^t) = y + [2t+2, 5t+4] = y + 2\lceil \frac{k}{3} \rceil + [0, k] = L, \text{ where } k = 3t+2.$$

If $L = y + 2k + 2 \cdot [0, k] \in \mathcal{L}_3$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N}$, then we set $A = 0^y g^{4k}(-g)^{4k}$ and hence $L(A) = L$.

If $L = y + 2t + 2 + (\{0\} \cup [2, 2t+3]) \in \mathcal{L}_4$ with $y, t \in \mathbb{N}_0$, then we set $A = 0^y U_1(-U_1) g^{4t}(-g)^{4t}$ and obtain that $L(A) = y + 2t + 2 + (\{0\} \cup [2, 2t+3]) = L$ by Equation (4).

Finally we suppose that $L = y + k + ([0, k-2] \cup \{k\}) \in \mathcal{L}_5$ with $k \geq 3$ and $y \in \mathbb{N}_0$, and we distinguish two subcases. If $k = 2t$ with $t \geq 2$, then we set $A = 0^y U_5(-U_5) g^{4t-4}(-g)^{4t-4}$ and hence $L(A) = y + k + ([0, k-2] \cup \{k\}) = L$ by Equation (10). If $k = 2t+1$ with $t \geq 1$, then we set $A = 0^y U_5^2(-g)^4 g^{4t-4}(-g)^{4t-4}$ and hence $L(A) = y + k + ([0, k-2] \cup \{k\}) = L$ by Equation (9).

Step 2. We prove that for every $A \in \mathcal{B}(G^\bullet)$, $L(A) \in \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5$.

Let $A \in \mathcal{B}(G^\bullet)$. We may suppose that $\Delta(L(A)) \neq \emptyset$. By Lemma 4.7.1 we have to distinguish two cases.

CASE 1: $\Delta(L(A)) = \{1\}$.

Then $L(A)$ is an interval, say $L(A) = [y, y+k] = y + [0, k]$ with $y \geq 2$ and $k \geq 1$. If $k = 3$ and $y = 2$, then $L(A) = [2, 5]$, a contradiction to Lemma 4.7.1. Thus $k = 3$ implies that $y \geq 3$ and hence $L(A) \in \mathcal{L}_2$. If $k \leq 2$, then obviously $L(A) \in \mathcal{L}_2$. Suppose that $k \geq 4$. If $y = 2t$ with $t \geq 2$, then $y+k \leq 5t$ and hence $y = 2t \geq 2\lceil \frac{k}{3} \rceil$ which implies that $L(A) \in \mathcal{L}_2$. Suppose that $y = 2t+1$ with $t \in \mathbb{N}$. If $y+k \leq 5t+1$, then $y = 2t+1 \geq 1 + 2\lceil \frac{k}{3} \rceil$ which implies that $L(A) \in \mathcal{L}_2$. Otherwise $y+k = 5t+2$ and hence $L(A) = [2t+1, 5t+2] \in \mathcal{L}_2$.

CASE 2: $2 \in \Delta(L(A)) \subset [1, 2]$.

We freely use the classification of minimal zero-sum sequence given in Lemma 4.6. Since $2 \in \Delta(L(A))$, there are $k \in \mathbb{N}$ and $U_1, \dots, U_k, V_1, \dots, V_{k+2} \in \mathcal{A}(G)$ with $|U_1| \geq |U_2| \geq \dots \geq |U_k|$ such that

$$A = U_1 \cdot \dots \cdot U_k = V_1 \cdot \dots \cdot V_{k+2} \quad \text{and} \quad k+1 \notin L(A),$$

and we may suppose that k is minimal with this property. Then $[\min L(A), k] \in L(A)$ and there exists $k_0 \in [2, k]$ such that $|U_i| \geq 3$ for every $i \in [1, k_0]$ and $|U_i| = 2$ for every $i \in [k_0+1, k]$. We continue with two simple assertions.

A1. For each two distinct $i, j \in [1, k_0]$, we have that $3 \notin L(U_i U_j)$.

A2. $|\mathcal{L}(U_1 \cdot \dots \cdot U_{k_0})| \geq 2$.

Proof of A1. Assume to the contrary that there exist distinct $i, j \in [1, k_0]$ such that $3 \in \mathcal{L}(U_i U_j)$. This implies that $k+1 \in \mathcal{L}(A)$, a contradiction. \square

[Proof of A1]

Proof of A2. Assume to the contrary that $|\mathcal{L}(U_1 \cdot \dots \cdot U_{k_0})| = 1$. Then Lemma 4.2.2 implies that $\max \mathcal{L}(A) = \max \mathcal{L}(U_1 \cdot \dots \cdot U_{k_0}) + k - k_0 = k$, a contradiction. \square

[Proof of A2]

We use **A1** and **A2** without further mention and freely use Lemma 4.6 together with all its notation. We distinguish six subcases.

CASE 2.1: $U_1 \in S_5$.

Without loss of generality, we may assume that $U_1 = eg^3(e+g)$. We choose $j \in [2, k_0]$ and start with some preliminary observations. If $|U_j| = 5$, then the fact that $3 \notin \mathcal{L}(U_1 U_j)$ implies that $U_j = -U_1$. If $|U_j| = 4$, then $3 \notin \mathcal{L}(U_1 U_j)$ implies that $U_j \in \{g^2(e+g)^2, g^4, (-g)^4, (e+g)^4\}$. If $|U_j| = 3$, then $3 \notin \mathcal{L}(U_1 U_j)$ implies that $U_j \in \{(e+2g)g(e+g), g^2(2g), (e+g)^2(2g)\}$.

Now we distinguish three cases.

Suppose that $|U_2| = 5$. Then $U_2 = -U_1$ and by symmetry we obtain that $U_j \in \{g^4, (-g)^4\}$ for every $j \in [3, k_0]$. Let $i \in [k_0+1, k]$. If $U_i \neq e^2$, then $4 \in U_1 U_2 U_i$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. Therefore we obtain that

$$A = U_1(-U_1)(g^4)^{k_1}((-g)^4)^{k_2}(e^2)^{k_3} \quad \text{where } k_1, k_2, k_3 \in \mathbb{N}_0,$$

and without loss of generality we may assume that $k_1 \geq k_2$. Then it follows that $\mathcal{L}(A)$ is equal to

$$k_1 - k_2 + k_3 + \mathcal{L}(U_1(-U_1)(g^4)^{k_2}((-g)^4)^{k_2}) = k_3 + k_1 - k_2 + 2k_2 + 2 + (\{0\} \cup [2, 2k_2 + 3]),$$

which is an element of \mathcal{L}_4 .

Suppose that $|U_2| = 4$ and there exists $j \in [2, k_0]$ such that $U_j = (-g)^4$, say $j = 2$. Let $i \in [3, k_0]$. If $U_i \in \{g^2(e+g)^2, g^2(2g)\}$, then $3 \in \mathcal{L}(U_2 U_i)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. If $U_i \in \{(e+g)^4, (e+g)^2(2g), (e+2g)g(e+g)\}$, then $4 \in \mathcal{L}(U_1 U_2 U_i)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. Therefore $U_i \in \{g^4, (-g)^4\}$. Let $\tau \in [k_0+1, k]$. If $U_\tau \in \{(e+2g)^2, (2g)^2, (e+g)(e-g)\}$, then $4 \in \mathcal{L}(U_1 U_2 U_\tau)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. Therefore $U_\tau \in \{e^2, g(-g)\}$. Therefore we obtain that

$$A = U_1(g^4)^{k_1}((-g)^4)^{k_2}(g(-g))^{k_3}(e^2)^{k_4} \quad \text{where } k_1, k_3, k_4 \in \mathbb{N}_0 \text{ and } k_2 \in \mathbb{N}$$

and hence $\mathcal{L}(A)$ is equal to

$$\mathcal{L}((g^4)^{k_1+1}((-g)^4)^{k_2}(g(-g))^{k_3}(e^2)^{k_4}) = k_4 + \mathcal{L}(g^{4k_1+4+k_3}(-g)^{4k_2+k_3})$$

which is in \mathcal{L}_3 .

Suppose that $|U_2| \leq 4$ and for every $j \in [2, k_0]$, we have $U_j \neq (-g)^4$. Then $U_j \in \{g^2(e+g)^2, g^4, (e+g)^4, (e+2g)g(e+g), g^2(2g), (e+g)^2(2g)\}$. Since $\text{supp}(U_1 \cdot \dots \cdot$

$U_{k_0}) \subset \{e, g, 2g, e+g, e+2g\}$ and $v_e(U_1 \cdot \dots \cdot U_{k_0}) = 1$, Lemma 4.7.3 implies that $|\mathcal{L}(U_1 \cdot \dots \cdot U_{k_0})| = 1$, a contradiction.

CASE 2.2: $U_1 \in S_4^4$.

Without loss of generality, we may assume that $U_1 = eg(2g)(e+g)$. Let $j \in [2, k_0]$.

Suppose that $|U_j| = 4$. Since $3 \notin \mathcal{L}(U_1 U_j)$, we obtain that $U_j \in \{g^2(e+g)^2, g^4, (e+g)^4\}$. Thus $U_1 U_j = W_1 W_2$ with $|W_1| = 5$, where W_1, W_2 are atoms and hence we are back to CASE 2.1.

Suppose that $|U_j| = 3$. Since $3 \notin \mathcal{L}(U_1 U_j)$, we obtain that $U_j \in \{(e+2g)g(e+g), g^2(2g), (e+g)^2(2g)\}$. If $U_j \in \{g^2(2g), (e+g)^2(2g)\}$, then $U_1 U_j = W_1 W_2$ with $|W_1| = 5$, where W_1, W_2 are atoms and hence we are back to CASE 2.1. Thus it remains to consider the case where $U_j = (e+2g)g(e+g)$.

Therefore we have

$$U_1 \cdot \dots \cdot U_{k_0} = U_1 ((e+2g)g(e+g))^{k_1} \quad \text{where } k_1 \in \mathbb{N}.$$

Since $\text{supp}(U_1 \cdot \dots \cdot U_{k_0}) \subset \{e, g, 2g, e+g, e+2g\}$ and $v_e(U_1 \cdot \dots \cdot U_{k_0}) = 1$, Lemma 4.7.3 implies that $|\mathcal{L}(U_1 \cdot \dots \cdot U_{k_0})| = 1$, a contradiction.

CASE 2.3: $U_1 \in S_4^3$ and for every $i \in [2, k_0]$, we have $U_i \notin S_4^4$.

Without loss of generality, we may assume that $U_1 = eg^2(e+2g)$. Let $j \in [2, k_0]$.

Suppose that $|U_j| = 4$. Since $3 \notin \mathcal{L}(U_1 U_j)$, we obtain that $U_j \in \{-U_1, g^2(e+g)^2, g^2(e-g)^2, (e+g)^4, (e-g)^4, g^4\}$. If $U_j \in \{g^2(e+g)^2, g^2(e-g)^2, (e+g)^4, (e-g)^4\}$, then $U_1 U_j = W_1 W_2$ with $|W_1| = 5$, where W_1, W_2 are atoms and hence we are back to CASE 2.1. Thus it remains to consider the cases where $U_j = -U_1$ or $U_j = g^4$.

Suppose that $|U_j| = 3$. Since $3 \notin \mathcal{L}(U_1 U_j)$, we obtain that $U_j \in \{eg(e-g), (e+2g)g(e+g), g^2(2g), (e+g)^2(2g), (e-g)^2(2g)\}$. If $U_j \in \{eg(e-g), (e+2g)g(e+g)\}$, then $U_1 U_j = W_1 W_2$ with $|W_1| = 5$, where W_1, W_2 are atoms and hence we are back to CASE 2.1. If $U_j \in \{(e+g)^2(2g), (e-g)^2(2g)\}$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^4$, where W_1, W_2 are atoms and hence we are back to CASE 2.2. Thus it remains to consider the case where $U_j = g^2(2g)$.

If $U_i \neq -U_1$ for every $i \in [2, k_0]$, then $U_1 \cdot \dots \cdot U_{k_0} = U_1 (g^4)^{k_1} (g^2(2g))^{k_2}$ where $k_1, k_2 \in \mathbb{N}_0$. Since $\text{supp}(U_1 \cdot \dots \cdot U_{k_0}) \subset \{e, g, 2g, e+g, e+2g\}$ and $v_e(U_1 \cdot \dots \cdot U_{k_0}) = 1$, Lemma 4.7.3 implies that $|\mathcal{L}(U_1 \cdot \dots \cdot U_{k_0})| = 1$, a contradiction. Thus there exists some $i \in [2, k_0]$, say $i = 2$, such that $U_2 = -U_1$. By symmetry we obtain that $k_0 = 2$. Let $\tau \in [3, k]$. If $U_\tau \in \{(2g)^2, (e+g)(e-g)\}$, then $4 \in \mathcal{L}(U_1 U_2 U_\tau)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. Therefore $A = U_1 (-U_1) (e^2)^{k_1} ((e+2g)^2)^{k_2} (g(-g))^{k_3}$ where $k_1, k_2, k_3 \in \mathbb{N}_0$. Since $[\min \mathcal{L}(A), 2+k_1+k_2+k_3] \subset \mathcal{L}(A)$, we obtain that $\mathcal{L}(A) = [\min \mathcal{L}(A), 2+y] \cup \{4+y\}$ where $y = k_1+k_2+k_3 \in \mathbb{N}_0$. For every atom V dividing A , we have that $|V| = 2$ or $|V| = 4$. Thus $\min \mathcal{L}(A) \geq 2 + \frac{y}{2}$ which implies that $\mathcal{L}(A) \in \mathcal{L}_5$.

CASE 2.4: $U_1 \in S_4^2$ and for every $i \in [2, k_0]$, we have $U_i \notin S_4^4 \cup S_4^3$.

Without loss of generality, we may assume that $U_1 = g^2(e+g)^2$. Let $j \in [2, k_0]$.

Suppose that $|U_j| = 4$. If $U_j \in \{g^2(e-g)^2, (-g)^2(e+g)^2, (-g)^4, (e-g)^4\}$, then $3 \in \mathcal{L}(U_1 U_j)$, a contradiction. Thus $U_j \in \{U_1, -U_1, g^4, (e+g)^4\}$.

Suppose that $|U_j| = 3$. If $U_j \in \{(e+2g)(-g)(e-g), (-g)^2(2g), (e-g)^2(2g)\}$, then $3 \in L(U_1U_j)$, a contradiction. If $U_j \in \{eg(e-g), e(-g)(e+g)\}$, then $U_1U_j = W_1W_2$ with $|W_1| = 5$, where W_1, W_2 are atoms and hence we are back to CASE 2.1. If $U_j = e(2g)(e+2g)$, then $U_1U_j = (e(e+g)g(2g))(g(e+g)(e+2g))$ and $e(e+g)g(e+2g) \in S_4^4$, going back to CASE 2.2. Thus it remains to consider the case where $U_j = g^2(2g)$ or $U_j = (e+g)^2(2g)$.

If $U_i \neq -U_1$ for every $i \in [2, k_0]$, then $\text{supp}(U_1 \cdots U_{k_0}) \subset \{g, 2g, e+g, e+2g\}$ is half-factorial by Lemma 4.7.3, a contradiction. Thus there exists some $i \in [2, k_0]$, say $i = 2$, such that $U_2 = -U_1$. By symmetry we obtain that $\{U_1, \dots, U_{k_0}\} = \{U_1, -U_1\}$. Let $\tau \in [k_0 + 1, k]$. If $U_\tau \in \{e^2, (2g)^2, (e+2g)^2\}$, then $4 \in L(U_1U_2U_\tau)$ and $k+1 \in L(U_1U_2U_\tau)$, a contradiction. Therefore $A = U_1^{k_1}(-U_1)^{k_2}(g(-g))^{k_3}((e+g)(e-g))^{k_4}$ where $k_1, k_2 \in \mathbb{N}$ and $k_3, k_4 \in \mathbb{N}_0$. If $k_1 + k_2 \geq 3$, by symmetry we assume that $k_1 \geq 2$, then $U_1^2(-U_1) = g^4(-g)^2(e+g)^2(e+g)(e-g)(e+g)(e-g)$ and hence $4 \in L(U_1^2(-U_1))$ which implies that $k+1 \in L(A)$, a contradiction. Thus $k_1 = k_2 = 1$ and hence $A = U_1(-U_1)(g(-g))^{k_3}((e+g)(e-g))^{k_4}$ where $k_3, k_4 \in \mathbb{N}_0$. Since $[\min L(A), 2+k_3+k_4] \in L(A)$, we obtain that $L(A) = [\min L(A), 2+y] \cup \{4+y\}$ where $y = k_3 + k_4 \in \mathbb{N}_0$. For every atom V dividing A , we have that $|V| = 2$ or $|V| = 4$. Thus $\min L(A) \geq 2 + \frac{y}{2}$ which implies that $L(A) \in \mathcal{L}_5$.

CASE 2.5: $U_1 \in S_4^1$ and for every $i \in [2, k_0]$, we have $U_i \notin S_4^4 \cup S_4^3 \cup S_4^2$.

Without loss of generality, we may assume that $U_1 = g^4$. Let $j \in [2, k_0]$.

Suppose that $|U_j| = 4$. If $U_j \in \{(e+g)^4, (e-g)^4\}$, then $U_1U_j = W_1W_2$ with $W_1 \in S_4^2$, where W_1, W_2 are atoms and hence we are back to CASE 2.4. Thus it remains to consider the case where $U_j = U_1$ or $U_j = -U_1$.

Suppose that $|U_j| = 3$. If $U_j \in \{(-g)^2(2g)\}$, then $3 \in L(U_1U_j)$, a contradiction. If $U_j \in \{e(-g)(e+g), (e+2g)(-g)(e-g)\}$, then $U_1U_j = W_1W_2$ with $|W_1| = 5$, where W_1, W_2 are atoms and hence we are back to CASE 2.1. If $U_j \in \{(e+g)^2(2g), (e-g)^2(2g)\}$, then $U_1U_j = W_1W_2$ with $W_1 \in S_4^2$, where W_1, W_2 are atoms and hence we are back to CASE 2.4. If $U_j = e(2g)(e+2g)$, then $U_1U_j = W_1W_2$ with $W_1 \in S_4^3$, where W_1, W_2 are atoms and hence we are back to CASE 2.3. Thus it remains to consider the case where $U_j = g^2(2g)$, or $U_j = eg(e-g)$, or $U_j = (e+2g)g(e+g)$.

First, suppose that $U_i \neq -U_1$ for every $i \in [2, k_0]$. Then

$$U_1 \cdots U_{k_0} = U_1^{k_1}(eg(e-g))^{k_2}((e+2g)g(e+g))^{k_3}(g^2(2g))^{k_4},$$

where $k_1 \in \mathbb{N}$ and $k_2, k_3, k_4 \in \mathbb{N}_0$. If $k_2 \geq 1$ and $k_3 \geq 1$, then $eg(e-g)(e+2g)g(e+g) = eg^2(e+2g)(e+g)(e-g)$, $eg^2(e+2g) \in S_4^3$ and hence we are back to CASE 2.3. Thus we may assume that $k_2 = 0$ or $k_3 = 0$. Since $\{g, 2g, e+g, e+2g\}$ and $\{g, 2g, e, e-g\}$ are both half-factorial by Lemma 4.7.3, we obtain that $|L(U_1 \cdots U_{k_0})| = 1$, a contradiction.

Second, suppose that there exists some $i \in [2, k_0]$, say $i = 2$, such that $U_2 = -U_1$. By symmetry we obtain that $\{U_1, \dots, U_{k_0}\} = \{U_1, -U_1\}$. Since $4 \in L(U_1 \cdot U_2 \cdot (2g)^2)$, $5 \in L(U_1U_2e^2(e-g)(e+g))$, and $5 \in L(U_1U_2(e+2g)^2(e-g)(e+g))$, we obtain that

$$\{U_{k_0+1}, \dots, U_k\} \subset \{(e+g)(e-g), g(-g)\} \quad \text{or}$$

$$\{U_{k_0+1}, \dots, U_k\} \subset \{e^2, (e+2g)^2, g(-g)\}.$$

This implies that

$$A = (g^4)^{k_1}((-g)^4)^{k_2}((e+g)(e-g))^{k_3}(g(-g))^{k_4} \quad \text{or}$$

$$A = (g^4)^{k_1}((-g)^4)^{k_2}(e^2)^{k_3}((e+2g)^2)^{k_4}(g(-g))^{k_5},$$

where $k_1, k_2 \in \mathbb{N}$ and $k_3, k_4, k_5 \in \mathbb{N}_0$.

Suppose that $A = (g^4)^{k_1}((-g)^4)^{k_2}((e+g)(e-g))^{k_3}(g(-g))^{k_4}$, where $k_1, k_2 \in \mathbb{N}$ and $k_3, k_4, k_5 \in \mathbb{N}_0$. If $k_1 \geq 2$ and $k_3 \geq 2$, then $g^4 g^4 (-g)^4 (e+g)(e-g)(e+g)(e-g) = (g(-g))^4 g^2 (e+g)^2 g^2 (e-g)^2$ and hence $6 \in L(g^4 g^4 (-g)^4 (e+g)(e-g)(e+g)(e-g))$. Thus $k+1 \in L(A)$, a contradiction. Therefore by symmetry $k_3 = 1$ or $k_1 = k_2 = 1$. If $k_3 = 1$, then $L(A) = 1 + L((g^4)^{k_1}((-g)^4)^{k_2}(g(-g))^{k_4}) \in \mathcal{L}_3$. If $k_1 = k_2 = 1$, then $L(A) = [\min L(A), 2+y] \cup \{4+y\}$ where $y = k_3 + k_4 \in \mathbb{N}_0$. For every atom V dividing A , we have that $|V| = 2$ or $|V| = 4$. Thus $\min L(A) \geq 2 + \frac{y}{2}$ which implies that $L(A) \in \mathcal{L}_5$.

Suppose that $A = (g^4)^{k_1}((-g)^4)^{k_2}(e^2)^{k_3}((e+2g)^2)^{k_4}(g(-g))^{k_5}$, where $k_1, k_2 \in \mathbb{N}$ and $k_3, k_4, k_5 \in \mathbb{N}_0$. If $k_1 \geq 2$, $k_3 \geq 1$, and $k_4 \geq 1$, then $g^4 g^4 (-g)^4 e^2 (e+2g)^2 = (g(-g))^4 (e(e+2g)g^2)^2$ and hence $6 \in L(g^4 g^4 (-g)^4 e^2 (e+2g)^2)$. Thus $k+1 \in L(A)$, a contradiction. Therefore by symmetry $k_3 = 0$, or $k_4 = 0$, or $k_1 = k_2 = 1$. If $k_3 = 0$ or $k_4 = 0$, then $L(A) = k_3 + k_4 + L((g^4)^{k_1}((-g)^4)^{k_2}(g(-g))^{k_5}) \in \mathcal{L}_3$. If $k_1 = k_2 = 1$, then $L(A) = [\min L(A), 2+y] \cup \{4+y\}$ where $y = k_3 + k_4 + k_5 \in \mathbb{N}_0$. For every atom V dividing A , we have that $|V| = 2$ or 4 . Thus $\min L(A) \geq 2 + \frac{y}{2}$ which implies that $L(A) \in \mathcal{L}_5$.

CASE 2.6: $|U_1| = 3$.

Let $j \in [2, k_0]$. We distinguish three subcases.

First, we suppose that $U_1 \in S_3^3$, and without restriction we may assume that $U_1 = eg(e-g)$. If $U_j = -U_1$, then $3 \in L(U_1 U_j)$, a contradiction. If $U_j \in \{(-g)^2(2g), (e+g)^2(2g), e(2g)(e+2g)\}$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^4$ where W_1, W_2 are atoms and hence we are back to CASE 2.2. If $U_j \in \{(e+2g)g(e+g), (e+2g)(-g)(e-g)\}$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^3$ where W_1, W_2 are atoms and hence we are back to CASE 2.3. If $U_j = U_1$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^2$ where W_1, W_2 are atoms and hence we are back to CASE 2.4. Thus it remains to consider the case where $U_j = g^2(2g)$ or $(e-g)^2(2g)$. Then $U_1 \cdot \dots \cdot U_{k_0} = U_1 (g^2(2g))^{k_1} ((e-g)^2(2g))^{k_2}$ where $k_1, k_2 \in \mathbb{N}_0$. Since $\{e, g, 2g, e-g\}$ is half-factorial by Lemma 4.7.3, we obtain that $|L(U_1 \cdot \dots \cdot U_{k_0})| = 1$, a contradiction.

Second, we suppose that $U_1 \in S_3^2$, and without restriction we may assume that $U_1 = g^2(2g)$ and $U_j \notin S_3^3$. If $U_j = -U_1$, then $3 \in L(U_1 U_j)$. If $U_j = U_1$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^1$ where W_1, W_2 are atoms and hence we are back to CASE 2.5. If $U_j \in \{(e+g)^2(2g), (e-g)^2(2g)\}$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^2$ where W_1, W_2 are atoms and hence we are back to CASE 2.4. If $U_j = e(2g)(e+2g)$, then $U_1 U_j = W_1 W_2$ with $W_1 \in S_4^3$ where W_1, W_2 are atoms and hence we are back to CASE 2.3.

Third, we suppose that $U_1 \in S_3^1$, and without restriction we assume that $U_j \in S_3^1$. Thus $3 \in L(U_1 U_j)$, a contradiction. \square

4.3 The system of sets of lengths of C_2^4

Now we give a complete description of the system of sets of lengths of C_2^4 .

Theorem 4.8. $\mathcal{L}(C_2^4) = \mathcal{L}_1 \cup \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5 \cup \mathcal{L}_6 \cup \mathcal{L}_7 \cup \mathcal{L}_8$, where

$$\begin{aligned}
\mathcal{L}_1 &= \{\{y\} \mid y \in \mathbb{N}_0\}, \\
\mathcal{L}_2 &= \{y + 2k + 3 \cdot [0, k] \mid y, k \in \mathbb{N}_0\}, \\
\mathcal{L}_3 &= \{y + [2, 3], y + [2, 4], y + [3, 6], y + [3, 7], y + [4, 9] \mid y \in \mathbb{N}_0\} \cup \\
&\quad \{y + [m, m+k] \mid y \in \mathbb{N}_0, k \geq 6, m \text{ minimal with } m+k \leq 5m/2\} \\
&= \{y + \left\lceil \frac{2k}{3} \right\rceil + [0, k] \mid y \in \mathbb{N}_0, k \in \mathbb{N} \setminus \{1, 3\}\} \cup \\
&\quad \{y + 3 + [0, 3], y + 2 + [0, 1] \mid y \in \mathbb{N}_0\}, \\
\mathcal{L}_4 &= \{y + 2k + 2 \cdot [0, k] \mid y, k \in \mathbb{N}_0\}, \\
\mathcal{L}_5 &= \{y + k + 2 + ([0, k] \cup \{k+2\}) \mid y \in \mathbb{N}_0, k \in \mathbb{N}\}, \\
\mathcal{L}_6 &= \{y + 2 \left\lceil \frac{k}{3} \right\rceil + 2 + (\{0\} \cup [2, k+2]) \mid y \in \mathbb{N}_0, k \geq 5 \text{ or } k = 3\}, \\
\mathcal{L}_7 &= \{y + 2k + 3 + \{0, 1, 3\} + 3 \cdot [0, k] \mid y, k \in \mathbb{N}_0\} \cup \\
&\quad \{y + 2k + 4 + \{0, 1, 3\} + 3 \cdot [0, k] \cup \{y + 5k + 8\} \mid y, k \in \mathbb{N}_0\}, \text{ and} \\
\mathcal{L}_8 &= \{y + 2k + 3 + \{0, 2, 3\} + 3 \cdot [0, k] \mid y, k \in \mathbb{N}_0\} \cup \\
&\quad \{y + 2k + 4 + \{0, 2, 3\} + 3 \cdot [0, k] \cup \{y + 5k + 9\} \mid y, k \in \mathbb{N}_0\}.
\end{aligned}$$

We note that the system of sets of lengths of C_2^4 is richer than that of the other groups we considered. A reason for this is that the set $\Delta^*(C_2^4)$ is largest, namely $\{1, 2, 3\}$ (this fact was also crucial in the proof of Theorem 3.5). We recall some useful facts in the lemma below.

Lemma 4.9. Let $G = C_2^4$, and let $A \in \mathcal{B}(G)$.

1. $\Delta(G) = [1, 3]$, and if $3 \in \Delta(L(A))$, then $\Delta(L(A)) = \{3\}$ and there is a basis (e_1, \dots, e_4) of G such that $\text{supp}(A) \setminus \{0\} = \{e_1, \dots, e_4, e_1 + \dots + e_4\}$.
2. $\rho_{2k+1}(G) = 5k+2$ for all $k \in \mathbb{N}$.

Proof. 1. The first statement follows from [20, Theorem 6.8.3], and the second statement from [25, Lemma 3.10].

2. See [20, Theorem 6.3.4]. □

In the following result we characterize which intervals are sets of lengths for C_2^4 . It turns out that, with a single exception, the sole restriction is the one implied by elasticity.

Proposition 4.10. *Let $G = C_2^4$ and let $2 \leq l_1 \leq l_2$ be integers. Then $[l_1, l_2] \in \mathcal{L}(G)$ if and only if $l_2/l_1 \leq 5/2$ and $(l_1, l_2) \neq (2, 5)$.*

Proof. Suppose that $[l_1, l_2] \in \mathcal{L}(G)$ with integers $2 \leq l_1 \leq l_2$. Then (2) implies that $l_2/l_1 \leq \rho(G) = 5/2$. Moreover, $[2, 5] = [2, D(G)] \notin \mathcal{L}(G)$ by [20, Theorem 6.6.3].

Conversely, we need to show that for integers $2 \leq l_1 \leq l_2$ with $(l_1, l_2) \neq (2, 5)$ and $l_2/l_1 \leq 5/2$, we have $[l_1, l_2] \in \mathcal{L}(G)$. We start with an observation that reduces the problem to constructing these sets of intervals for extremal choices of the endpoints.

Let $k \in \mathbb{N}$. If $m \in \mathbb{N}$ such that $[m, m+k] \in \mathcal{L}(G)$, then $y + [m, m+k] \in \mathcal{L}(G)$ for all $y \in \mathbb{N}_0$. Thus let $m_k = \max\{2, \lceil \frac{2k}{3} \rceil\}$ if $k \in \mathbb{N} \setminus \{3\}$ and $m_3 = 3$. Therefore we only need to prove that $[m_k, m_k+k] \in \mathcal{L}(G)$.

For $k \in [1, 5]$ we are going to realize sets $[m_k, m_k+k]$ as sets of lengths. Then we handle the case $k \geq 6$.

If $k \in \{1, 3\}$, then the sets $[2, 3], [3, 6] \in \mathcal{L}(C_2^3) \subset \mathcal{L}(G)$. To handle the case $k = 2$, we have to show that $[2, 4] \in \mathcal{L}(G)$. Let (e_1, \dots, e_4) be a basis of G and $e_0 = e_1 + \dots + e_4$. If

$$U_1 = e_0 \cdot \dots \cdot e_4 \quad \text{and} \quad U_2 = e_1 e_2 (e_1 + e_3) (e_2 + e_4) (e_3 + e_4),$$

then $\max L(U_1 U_2) < 5$, and

$$\begin{aligned} U_1 U_2 &= (e_0 e_1 e_2 (e_3 + e_4)) ((e_1 + e_3) e_1 e_3) ((e_2 + e_4) e_2 e_4) \\ &= (e_0 (e_1 + e_3) (e_2 + e_4)) (e_1^2) (e_2^2) ((e_3 + e_4) e_3 e_4), \end{aligned}$$

shows that $L(U_1 U_2) = [2, 4]$. It remains to verify the following assertions.

- A1.** $[3, 7] \in \mathcal{L}(G)$ (this settles the case $k = 4$).
- A2.** $[4, 9] \in \mathcal{L}(G)$ (this settles the case $k = 5$).
- A3.** Let $k \geq 6$. Then $[\lceil \frac{2k}{3} \rceil, \lceil \frac{2k}{3} \rceil + k] \in \mathcal{L}(G)$.

Proof of A1. Clearly,

$$\begin{aligned} U_1 &= e_0 \cdot \dots \cdot e_4, \quad U_2 = e_1 e_2 (e_1 + e_3) (e_2 + e_4) (e_3 + e_4), \quad \text{and} \\ U_3 &= (e_1 + e_3) (e_2 + e_4) e_3 e_4 (e_1 + e_2) \end{aligned}$$

are minimal zero-sum sequences of lengths 5. Since

$$\begin{aligned} U_1 U_2 U_3 &= (e_0 (e_1 + e_2) (e_3 + e_4)) (e_1^2) (e_2^2) (e_3^2) (e_4^2) ((e_1 + e_3)^2) ((e_2 + e_4)^2) \\ &= (e_0 (e_1 + e_2) (e_3 + e_4)) ((e_1 + e_3) e_1 e_3)^2 ((e_2 + e_4)^2) (e_2^2) (e_4^2) \\ &= (e_0 (e_1 + e_2) (e_3 + e_4)) ((e_1 + e_3) e_1 e_3)^2 ((e_2 + e_4) e_2 e_4)^2 \\ &= U_2 (e_0 (e_1 + e_2) (e_1 + e_3) e_1 e_4) ((e_2 + e_4) e_2 e_4) (e_3^2), \end{aligned}$$

it follows that $L(U_1 U_2 U_3) = [3, 7]$.

Proof of A2. We use the same notation as in A1, set $U_4 = (e_1 + e_2)(e_1 + e_3)(e_2 + e_4)(e_3 + e_4)$, and assert that $L(U_1^2 U_2 U_4) = [4, 9]$. Clearly, $4 \in L(U_1^2 U_2 U_4)$ and $\max L(U_1^2 U_2 U_4) < 10$. Since

$$\begin{aligned} U_1^2 U_2 U_4 &= (e_0 e_1 e_2 (e_3 + e_4)) ((e_1 + e_3) e_1 e_3) ((e_2 + e_4) e_2 e_4) U_1 U_4 \\ &= (e_0 (e_1 + e_3) (e_2 + e_4)) (e_1^2) (e_2^2) ((e_3 + e_4) e_3 e_4) U_1 U_4 \\ &= \prod_{v=0}^4 (e_v^2) U_2 U_4 \\ &= ((e_1 + e_3)^2) ((e_2 + e_4)^2) ((e_3 + e_4) e_3 e_4)^2 (e_0^2) (e_1^2) (e_2^2) ((e_1 + e_2) e_1 e_2) \\ &= ((e_1 + e_3)^2) ((e_2 + e_4)^2) ((e_3 + e_4)^2) (e_3^2) (e_4^2) (e_0^2) (e_1^2) (e_2^2) ((e_1 + e_2) e_1 e_2), \end{aligned}$$

the assertion follows.

Proof of A3. We proceed by induction on k . For $k = 6$, we have to verify that $[4, 10] \in \mathcal{L}(G)$. We use the same notation as in A1, and assert that $L(U_1^2 U_2^2) = [4, 10]$. Clearly, $\{4, 10\} \subset L(U_1^2 U_2^2) \subset [4, 10]$. Since

$$\begin{aligned} U_1^2 U_2^2 &= (e_0 e_1 e_2 (e_3 + e_4)) ((e_1 + e_3) e_1 e_3) ((e_2 + e_4) e_2 e_4) U_1 U_2 \\ &= (e_0 e_1 e_2 (e_3 + e_4))^2 ((e_1 + e_3) e_1 e_3)^2 ((e_2 + e_4) e_2 e_4)^2 \\ &= \prod_{v=0}^4 (e_v^2) U_2^2 \\ &= (e_0 (e_1 + e_3) (e_2 + e_4))^2 (e_1^2)^2 (e_2^2)^2 ((e_3 + e_4) e_3 e_4)^2 \\ &= ((e_1 + e_3)^2) ((e_2 + e_4)^2) ((e_3 + e_4) e_3 e_4)^2 (e_0^2) (e_1^2)^2 (e_2^2)^2 \end{aligned}$$

it follows that $[5, 9] \subset L(U_1^2 U_2^2)$, and hence $L(U_1^2 U_2^2) = [4, 10]$.

If $k = 7$, then $[5, 12] \supset L(U_1^3 U_2 U_3) \supset L(U_1 U_2 U_3) + L(U_1^2) = [3, 7] + \{2, 5\} = [5, 12]$ which implies that $[5, 12] \in \mathcal{L}(G)$. If $k = 8$, then $[6, 14] \supset L(U_1^4 U_2 U_4) \supset L(U_1^2 U_2 U_4) + L(U_1^2) = [4, 9] + \{2, 5\} = [6, 14]$ which implies that $[6, 14] \in \mathcal{L}(G)$. Suppose that $k \geq 9$, and that the assertion holds for all $k' \in [6, k-1]$. Then the set $[\lceil \frac{2(k-3)}{3} \rceil, \lceil \frac{2(k-3)}{3} \rceil + k-3] \in \mathcal{L}(G)$. This implies that $[\lceil \frac{2k}{3} \rceil, \lceil \frac{2k}{3} \rceil + k] = [\lceil \frac{2(k-3)}{3} \rceil, \lceil \frac{2(k-3)}{3} \rceil + k-3] + \{2, 5\} \in \mathcal{L}(G)$. \square

We now proceed to prove Theorem 4.8.

Proof (Theorem 4.8). Let (e_1, e_2, e_3, e_4) be a basis of $G = C_2^4$. We set $e_0 = e_1 + e_2 + e_3 + e_4$, $U = e_0 e_1 e_2 e_3 e_4$, and $V = e_1 e_2 e_3 (e_1 + e_2 + e_3)$.

Step 0. Some elementary constructions.

Let $t_1 \geq 2$, $t_2 \geq 2$, $t = t_1 + t_2$, and

$$L_{t_1, t_2} = \begin{cases} \{t\} \cup [t+2, 5\lfloor t_1/2 \rfloor + 4(t/2 - \lfloor t_1/2 \rfloor)] & \text{if } t \text{ is even,} \\ \{t\} \cup [t+2, 5\lfloor t_1/2 \rfloor + 4((t-1)/2 - \lfloor t_1/2 \rfloor) + 1] & \text{if } t \text{ is odd.} \end{cases}$$

Since $\mathsf{L}(U^2V^2) = \{4\} \cup [6, 9]$, we have that $\mathsf{L}(U^{t_1}V^{t_2}) \supset \mathsf{L}(U^2V^2) + \mathsf{L}(U^{t_1-2}V^{t_2-2}) = L_{t_1, t_2}$. Note that for every atom W dividing $U^{t_1}V^{t_2}$, we have

$$W = \begin{cases} U & \text{if } |W| = 5, \\ V & \text{if } |W| = 4, \\ e_0e_4(e_1 + e_2 + e_3) & \text{if } |W| = 3. \end{cases}$$

Assume to the contrary that $t+1 \in \mathsf{L}(U^{t_1}V^{t_2})$. Then there exist $t_3, t_4, t_5 \in \mathbb{N}_0$ and atoms $W_1, \dots, W_{t_3+t_4+1}$ such that $U^{t_3}V^{t_4} = W_1 \dots W_{t_3+t_4+1}$ with $t_3+t_4 \geq 2$, $t_5 \leq \min\{t_3, t_4\}$, $|W_i| = 3$ for $i \in [1, t_5]$, and $|W_i| = 2$ for $i \in [t_5+1, t_3+t_4+1]$. It follows that $5t_3 + 4t_4 = 3t_5 + 2(t_3+t_4+1-t_5) \leq 3t_3 + 2t_4 + 2$ and hence $t_3+t_4 \leq 1$, a contradiction. Therefore $t+1 \notin \mathsf{L}(U^{t_1}V^{t_2})$ and

$$\mathsf{L}(U^{t_1}V^{t_2}) = L_{t_1, t_2}. \quad (11)$$

Note that for every atom W dividing U^rV with $r \geq 2$ and $e_1 + e_2 + e_3 \mid W$, we have $W = V$ or $W = e_0e_4(e_1 + e_2 + e_3)$. It follows that for all $r \geq 2$

$$\begin{aligned} & \mathsf{L}(U^rV) \\ &= (1 + \mathsf{L}(U^r)) \cup (1 + \mathsf{L}(e_1^2e_2^2e_3^2U^{r-1})) \quad (12) \\ &= \begin{cases} r+1 + \{0, 2, 3\} + 3 \cdot [0, r/2-1], & \text{if } r \text{ is even,} \\ r+1 + \{0, 2, 3\} + 3 \cdot [0, (r-1)/2-1] \cup \{r+1 + (3r-3)/2+2\}, & \text{if } r \text{ is odd.} \end{cases} \end{aligned}$$

Note that for every atom W dividing $U^rVe_4^2e_0^2$ with $r \geq 2$ and $e_1 + e_2 + e_3 \mid W$, we have $W = V$ or $W = e_0e_4(e_1 + e_2 + e_3)$. It follows that for all $r \geq 2$

$$\begin{aligned} & \mathsf{L}(U^rVe_4^2e_0^2) \\ &= (1 + \mathsf{L}(U^r e_4^2 e_0^2)) \cup (1 + \mathsf{L}(U^{r+1})) \quad (13) \\ &= \begin{cases} r+2 + \{0, 1, 3\} + 3 \cdot [0, (r+1)/2-1], & \text{if } r \text{ is odd,} \\ r+2 + \{0, 1, 3\} + 3 \cdot [0, r/2-1] \cup \{r+2 + 3r/2+1\}, & \text{if } r \text{ is even.} \end{cases} \end{aligned}$$

Step 1. We prove that for every $L \in \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5 \cup \mathcal{L}_6 \cup \mathcal{L}_7 \cup \mathcal{L}_8$, there exists an $A \in \mathcal{B}(G)$ such that $L = \mathsf{L}(A)$. We distinguish seven cases.

If $L = y + 2k + 3 \cdot [0, k] \in \mathcal{L}_2$ with $y, k \in \mathbb{N}_0$, then $L = \mathsf{L}(0^y U^{2k}) \in \mathcal{L}(G)$.

If $L \in \mathcal{L}_3$, then the claim follows from Proposition 4.10.

If $L = y + 2k + 2 \cdot [0, k] \in \mathcal{L}_4$ with $y, k \in \mathbb{N}_0$, then Proposition 3.3.4 implies that $L \in \mathcal{L}(C_2^3) \subset \mathcal{L}(G)$.

Suppose that $L = y + k + 2 + ([0, k] \cup \{k+2\}) \in \mathcal{L}_5$ with $k \in \mathbb{N}$ and $y \in \mathbb{N}_0$. Note that $\mathsf{L}(V^2(e_1 + e_4)^2(e_2 + e_4)^2(e_3 + e_4)^2(e_1 + e_2 + e_3 + e_4)^2) = [4, 6] \cup \{8\}$. If

k is even, then we set $A = 0^y V^2 (e_1 + e_4)^k (e_2 + e_4)^k (e_3 + e_4)^k (e_1 + e_2 + e_3 + e_4)^k$ and obtain that $L(A) = L$ by Lemma 4.2.3. If k is odd, then we set $A = 0^y V^2 (e_1 + e_4)^{k+1} (e_2 + e_4)^{k+1} (e_3 + e_4)^{k-1} (e_1 + e_2 + e_3 + e_4)^{k-1}$ and obtain that $L(A) = L$ by Lemma 4.2.3.

Suppose that $L = y + 2\lceil \frac{k}{3} \rceil + 2 + (\{0\} \cup [2, k+2]) \in \mathcal{L}_6$ with $(k \geq 5 \text{ or } k = 3)$ and $y \in \mathbb{N}_0$. If $k \equiv 0 \pmod{3}$, then we set $A = 0^y U^{2k/3} V^2$ and hence $L(A) = L$ by Equation (11). If $k \equiv 2 \pmod{3}$, then we set $A = 0^y U^{(2k-4)/3} V^4$ and hence $L(A) = L$ by (11). If $k \equiv 1 \pmod{3}$, then we set $A = 0^y U^{(2k-8)/3} V^6$ and obtain that $L(A) = L$ by Equation (11).

Suppose that $L \in \mathcal{L}_7$. If $L = y + 2k + 3 + \{0, 1, 3\} + 3 \cdot [0, k]$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N}_0$, then we set $A = 0^y U^{2k+1} V e_4^2 (e_1 + e_2 + e_3 + e_4)^2$ and obtain that $L(A) = L$ by Equation (13). If $L = y + 2k + 4 + \{0, 1, 3\} + 3 \cdot [0, k] \cup \{y + 5k + 8\}$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N}_0$, then we set $A = 0^y U^{2k+2} V e_4^2 (e_1 + e_2 + e_3 + e_4)^2$ and obtain that $L(A) = L$ by Equation (13).

Suppose that $L \in \mathcal{L}_8$. If $L = y + 2k + 3 + \{0, 2, 3\} + 3 \cdot [0, k]$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N}_0$, then we set $A = 0^y U^{2k+2} V$ and hence $L(A) = L$ by Equation (12). If $L = y + 2k + 4 + \{0, 2, 3\} + 3 \cdot [0, k] \cup \{y + 5k + 9\}$ with $y \in \mathbb{N}_0$ and $k \in \mathbb{N}_0$, then we set $A = 0^y U^{2k+3} V e_4^2 (e_1 + e_2 + e_3 + e_4)^2$ and obtain that $L(A) = L$ by Equation (12).

Step 2. We prove that for every $A \in \mathcal{B}(G^\bullet)$, $L(A) \in \mathcal{L}_2 \cup \mathcal{L}_3 \cup \mathcal{L}_4 \cup \mathcal{L}_5 \cup \mathcal{L}_6 \cup \mathcal{L}_7 \cup \mathcal{L}_8$.

Let $A \in \mathcal{B}(G^\bullet)$. We may suppose that $\Delta(L(A)) \neq \emptyset$. By Lemma 4.9.1 we have to distinguish four cases.

CASE 1: $\Delta(L(A)) = \{3\}$.

By Lemma 4.9, there is a basis of G , say (e_1, e_2, e_3, e_4) , such that $\text{supp}(A) = \{e_1, \dots, e_4, e_0\}$. Let $n \in \mathbb{N}_0$ be maximal such that $U^{2n} \mid A$. Then there exist a proper subset $I \subset [0, 4]$, a tuple $(m_i)_{i \in I} \in \mathbb{N}_0^{(I)}$, and $\varepsilon \in \{0, 1\}$ such that

$$A = U^\varepsilon U^{2n} \prod_{i \in I} (e_i^2)^{m_i}.$$

Using [25, Lemma 3.6.1], we infer that

$$L(A) = \varepsilon + \sum_{i \in I} m_i + L(U^{2n}) = \varepsilon + \sum_{i \in I} m_i + (2n + 3 \cdot [0, n]) \in \mathcal{L}_2.$$

CASE 2: $\Delta(L(A)) = \{1\}$.

Then $L(A)$ is an interval, and it is a direct consequence of Proposition 4.10 that $L(A) \in \mathcal{L}_3$.

CASE 3: $\Delta(L(A)) = \{2\}$.

The following reformulation turns out to be convenient. Clearly, we have to show that for every $L \in \mathcal{L}(G)$ with $\Delta(L) = \{2\}$ there exist $y' \in \mathbb{N}_0$ and $k' \in \mathbb{N}$ such that $L = y' + 2 \cdot [k', 2k']$, which is equivalent to $\rho(L) = \max L / \min L \leq 2$. Assume to the contrary that there is an $L \in \mathcal{L}(G)$ with $\Delta(L) = \{2\}$ such that $\max L \geq 2 \min L + 1$. We choose one such $L \in \mathcal{L}(G)$ with $\min L$ being minimal, and we choose a $B \in \mathcal{B}(G)$ with $L(B) = L$. Since $\min L$ is minimal, we obtain that

$0 \nmid B$. Consequently, $|B| \geq 2 \max L \geq 4 \min L + 2$. Since $D(G) = 5$, it follows that a factorization of minimal length of B contains at least two (possibly equal) minimal zero-sum sequences U_1, U_2 with $|U_1| = |U_2| = 5$, say $U_1 = e_0 \dots e_4$.

If $U_1 = U_2$, then $5 \in L(U_1 U_2)$ and thus $\min L + 3 \in L$, contradicting the fact that $\Delta(L) = \{2\}$. Thus $U_1 \neq U_2$. We assert that $3 \in L(U_1 U_2)$, and thus obtain again a contradiction to the fact that $\Delta(L) = \{2\}$.

Let $g \in G$ with $g \mid U_2$ but $g \nmid U_1$. Then g is the sum of two elements from U_1 , say $g = e_1 + e_2$. Therefore $g(e_1 e_2)^{-1} U_1$ is a minimal zero-sum sequence, whereas the sequence $(e_1 e_2)g^{-1} U_2$ cannot be a minimal zero-sum sequence because it has length 6. Since $g^{-1} U_2$ is zero-sum free, every minimal zero-sum sequence dividing $(e_1 e_2)g^{-1} U_2$ must contain e_1 or e_2 . This shows that $L((e_1 e_2)g^{-1} U_2) = \{2\}$ and thus $3 \in L(U_1 U_2)$.

CASE 4: $\Delta(L(A)) = \{1, 2\}$.

Let $k \in L(A)$ be minimal such that A has a factorization of the form $A = U_1 \cdot \dots \cdot U_k = V_1 \cdot \dots \cdot V_{k+2}$, where $k+1 \notin L(A)$ and $U_1, \dots, U_k, V_1, \dots, V_{k+2} \in \mathcal{A}(G)$ with $|U_1| \geq |U_2| \geq \dots \geq |U_k|$. Without restriction we may suppose that the tuple

$$(|\{i \in [1, k] \mid |U_i| = 5\}|, |\{i \in [1, k] \mid |U_i| = 4\}|, |\{i \in [1, k] \mid |U_i| = 3\}|) \in \mathbb{N}_0^3 \quad (14)$$

is maximal (with respect to the lexicographic order) among all factorizations of A of length k . By definition of k , we have $[\min L(A), k] \in L(A)$. Let $k_0 \in [2, k]$ such that $|U_i| \geq 3$ for every $i \in [1, k_0]$ and $|U_i| = 2$ for every $i \in [k_0 + 1, k]$. We start with the following assertion.

A.

1. For each two distinct $i, j \in [1, k_0]$, we have $3 \notin L(U_i U_j)$.
2. For each two distinct $i, j \in [1, k_0]$ with $|U_i| = |U_j| = 5$, we have $U_i = U_j$.
3. For each two distinct $i, j \in [1, k_0]$ with $|U_i| = 5$ and $|U_j| = 4$, we have $|\gcd(U_i, U_j)| = 3$.
4. Let $i, j \in [1, k_0]$ be distinct with $|U_i| = |U_j| = 4$, say $U_i = f_1 f_2 f_3 (f_1 + f_2 + f_3)$ where (f_1, f_2, f_3, f_4) a basis of G . Then $U_j = U_i$, or $U_j = (f_1 + f_4)(f_2 + f_4)(f_3 + f_4)(f_1 + f_2 + f_3 + f_4)$, or $U_j = f_4(f_1 + f_2 + f_4)(f_2 + f_3 + f_4)(f_1 + f_3 + f_4)$. Furthermore, if $U_i \neq U_j$, then for all $t \in [1, k_0] \setminus \{i, j\}$, we have $|U_t| \neq 4$.
5. Let $i, j \in [1, k_0]$ be distinct with $|U_i| = 5$ and $|U_j| = 3$. Then there exist $g_1, g_2, g_3 \in G$ such that $g_1 g_2 g_3 \mid U_i$ and $U_j = (g_1 + g_2)(g_2 + g_3)(g_3 + g_1)$. Furthermore, for all $t \in [1, k_0] \setminus \{i, j\}$, we have $|U_t| = 3$.
6. Let $i, j \in [1, k_0]$ be distinct with $|U_i| = 4$ and $|U_j| = 3$. Then $|\gcd(U_i, U_j)| = 0$, and there exist $g, g_1, g_2 \in G$ such that $g \mid U_j$, $g_1 g_2 \mid U_i$ and $g = g_1 + g_2$. Furthermore, for all $t \in [1, k_0] \setminus \{i, j\}$, we have $|U_t| = 3$.
7. For each two distinct $i, j \in [1, k_0]$ with $|U_i| = |U_j| = 3$, we have $|\gcd(U_i, U_j)| = 0$.

Proof of A.

1. If there exist distinct $i, j \in [1, k_0]$ such that $3 \in L(U_i U_j)$, then $k+1 \in L(A)$, a contradiction.

2. Since $|U_i| = 5$ and $U_j \neq U_i$, there exist $g, g_1, g_2 \in G$ with $g \mid U_j$ and $g_1g_2 \mid U_i$ such that $g = g_1 + g_2$. Thus $U_i(g_1g_2)^{-1}g$ is an atom and $U_jg^{-1}g_1g_2$ is a product of two atoms which implies that $3 \in L(U_iU_j)$, a contradiction.

3. Since $|U_i| = 5$ and $U_j \neq U_i$, there exist $g, g_1, g_2 \in G$ with $g \mid U_j$ and $g_1g_2 \mid U_i$ such that $g = g_1 + g_2$. Thus gg_1g_2 is an atom and $U_iU_j(gg_1g_2)^{-1}$ is a sequence of length 6. By 1., $2 \notin L(U_iU_j(gg_1g_2)^{-1})$ which implies that $L(U_iU_j(gg_1g_2)^{-1}) = \{3\}$ and hence $|\gcd(U_i, U_j)| = 3$.

4. We set $G_1 = \langle f_1, f_2, f_3 \rangle$ and distinguish three cases.

Case (i): $U_j \in \mathcal{B}(G_1)$. Since $3 \notin L(U_iU_j)$, we obtain that $U_j = U_i$.

Case (ii): $U_j = (g_1 + f_4)(g_2 + f_4)g_3g_4$ with $g_1g_2g_3g_4 \in \mathcal{B}(G_1)$.

If $g_3, g_4 \in \{f_1, f_2, f_3, f_1 + f_2 + f_3\}$, then $3 \in L(U_iU_j)$, a contradiction. Thus, without loss of generality, we may assume that $g_3 = f_1 + f_2 \notin \{f_1, f_2, f_3, f_1 + f_2 + f_3\}$. Thus $g_3f_3(f_1 + f_2 + f_3)$ is an atom and $(g_1 + f_4)(g_2 + f_4)f_1f_2g_4$ is a zero-sum sequence of length 5. Since $3 \notin L(U_iU_j)$, we have that $(g_1 + f_4)(g_2 + f_4)f_1f_2g_4$ is an atom of length 5, a contradiction to the maximality condition in Equation (14).

Case (iii): $U_j = (g_1 + f_4)(g_2 + f_4)(g_3 + f_4)(g_4 + f_4)$ with $g_1g_2g_3g_4 \in \mathcal{B}(G_1)$.

First, suppose that $g_1g_2g_3g_4$ is an atom. If $g_1g_2g_3g_4 \neq U_i$, then there exist an element $h \in \{f_1, f_2, f_3, f_1 + f_2 + f_3\}$ and distinct $t_1, t_2 \in [1, 4]$, say $t_1 = 1, t_2 = 2$, such that $h = g_1 + g_2 = (g_1 + f_4) + (g_2 + f_4)$. Thus $U_ih^{-1}(g_1 + f_4)(g_2 + f_4)$ is a zero-sum sequence of length 5 and $h(g_3 + f_4)(g_4 + f_4)$ is an atom. It follows that $U_ih^{-1}(g_1 + f_4)(g_2 + f_4)$ is atom of length 5 since $3 \notin L(U_iU_j)$, a contradiction to the maximality condition in Equation (14). Therefore $g_1g_2g_3g_4 = U_i$ which implies that $U_j = (f_1 + f_4)(f_2 + f_4)(f_3 + f_4)(f_1 + f_2 + f_3 + f_4)$.

Second, suppose that $g_1g_2g_3g_4$ is not an atom. Without loss of generality, we may assume that $g_1 = 0$ and $g_2g_3g_4$ is an atom. If $\{g_2, g_3, g_4\} \cap \{f_1, f_2, f_3, f_1 + f_2 + f_3\} \neq \emptyset$, say $g_2 \in \{f_1, f_2, f_3, f_1 + f_2 + f_3\}$, then $g_2(g_3 + f_4)(g_4 + f_4)$ is an atom and $U_ig_2^{-1}f_4(g_2 + f_4)$ is a zero-sum sequence of length 5. It follows that $U_ig_2^{-1}f_4(g_2 + f_4)$ is atom of length 5 because $3 \notin L(U_iU_j)$, a contradiction to the maximality condition in Equation (14). Therefore $\{g_2, g_3, g_4\} \cap \{f_1, f_2, f_3, f_1 + f_2 + f_3\} = \emptyset$ which implies that $g_2g_3g_4 = (f_1 + f_2)(f_2 + f_3)(f_1 + f_3)$ and hence $U_j = f_4(f_1 + f_2 + f_4)(f_2 + f_3 + f_4)(f_1 + f_3 + f_4)$.

Now suppose that $U_i \neq U_j$, and assume to the contrary there exists a $t \in [1, k_0] \setminus \{i, j\}$ such that $|U_t| = 4$. If $U_t \notin \{U_i, U_j\}$, then $U_iU_jU_t = (f_1f_2f_3(f_1 + f_2 + f_3))((f_1 + f_4)(f_2 + f_4)(f_3 + f_4)(f_1 + f_2 + f_3 + f_4))(f_4(f_1 + f_2 + f_4)(f_2 + f_3 + f_4)(f_1 + f_3 + f_4)) = (f_1(f_2 + f_4)(f_1 + f_2 + f_4))(f_2(f_3 + f_4)(f_2 + f_3 + f_4))(f_3(f_1 + f_4)(f_1 + f_3 + f_4))(f_4(f_1 + f_2 + f_3)(f_1 + f_2 + f_3 + f_4))$. Thus $4 \in L(U_iU_jU_t)$ and hence $k + 1 \in L(A)$, a contradiction. If $U_t \in \{U_i, U_j\}$, then we still have that $4 \in L(U_iU_jU_t)$ and hence $k + 1 \in L(A)$, a contradiction.

5. Since $3 \notin L(U_iU_j)$, we obtain that $|\gcd(U_i, U_j)| = 0$. Every $h \in \text{supp}(U_j)$ is the sum of two distinct elements from $\text{supp}(U_i)$. Thus there exist $g_1, g_2, g_3 \in G$ with $g_1g_2g_3 \mid U_i$ such that $U_j = (g_1 + g_2)(g_2 + g_3)(g_3 + g_1)$. Now we choose an element $t \in [1, k_0] \setminus \{i, j\}$, and have to show that $|U_t| = 3$. If $|U_t| = 5$, then $U_t = U_i$ by 2. and hence $4 \in L(U_iU_tU_j)$ which implies that $k + 1 \in L(A)$, a contradiction. Suppose that $|U_t| = 4$ and let $U_i = g_1g_2g_3g_4g_5$, where $g_4, g_5 \in G$. Then $|\gcd(U_i, U_t)| = 3$ by 3. and

by symmetry we only need to consider $\text{supp}(U_t) \setminus \text{supp}(U_i) \subset \{g_1 + g_2, g_1 + g_4, g_4 + g_5\}$. All the three cases imply that $4 \in \mathcal{L}(U_i U_t U_j)$. It follows that that $k + 1 \in \mathcal{L}(A)$, a contradiction.

6. If $|\text{gcd}(U_i, U_j)| = 2$, then $3 \in \mathcal{L}(U_i U_j)$, a contradiction. If $|\text{gcd}(U_i, U_j)| = 1$, then $U_1 U_2 = W_1 W_2$ with $W_1, W_2 \in \mathcal{A}(G)$ and $|W_2| = 5$, a contradiction to the maximality condition in Equation (14). Thus we obtain that $|\text{gcd}(U_i, U_j)| = 0$. Let (f_1, f_2, f_3, f_4) be a basis and $U_i = f_1 f_2 f_3 (f_1 + f_2 + f_3)$. Since $|U_j| = 3$, there exists a $g \in \text{supp}(U_j)$ such that $g \in \langle f_1, f_2, f_3 \rangle$. Since $|\text{gcd}(U_i, U_j)| = 0$, there exist $g_1, g_2 \in G$ such that $g_1 g_2 \mid U_i$ and $g = g_1 + g_2$.

Now we choose an element $t \in [1, k_0] \setminus \{i, j\}$ and have to show that $|U_t| = 3$. Note that 5. implies that $|U_t| \neq 5$, and we assume to the contrary that $|U_t| = 4$. Without restriction we may assume that $g = f_1 + f_2$, and by 4., we distinguish three cases. If $U_t = U_i$, then $f_1^2, f_2^2, g U_i (f_1 f_2)^{-1}, U_t (f_1 f_2)^{-1} U_j g^{-1}$ are atoms and hence $4 \in \mathcal{L}(U_i U_t U_j)$ which implies that $k + 1 \in \mathcal{L}(A)$, a contradiction. If $U_t = (f_1 + f_4)(f_2 + f_4)(f_3 + f_4)(f_1 + f_2 + f_3 + f_4)$, then $g(f_1 + f_2 + f_3)(f_1 + f_2 + f_3 + f_4)(f_1 + f_4)f_2$ is an atom of length 5 dividing $U_i U_j U_t$ and $U_i U_j U_t (g(f_1 + f_2 + f_3)(f_1 + f_2 + f_3 + f_4)(f_1 + f_4)f_2)^{-1}$ is a product of two atoms, a contradiction to the maximality condition in Equation (14). If $U_t = f_4(f_1 + f_2 + f_4)(f_2 + f_3 + f_4)(f_1 + f_3 + f_4)$, then $g f_2 f_3 f_4 (f_1 + f_3 + f_4)$ is an atom of length 5 dividing $U_i U_j U_t$ and $U_i U_j U_t (g f_2 f_3 f_4 (f_1 + f_3 + f_4))^{-1}$ is a product of two atoms, a contradiction to the maximality condition in Equation (14).

7. If $|\text{gcd}(U_i, U_j)| \geq 2$, then $U_i = U_j$ and hence $3 \in \mathcal{L}(U_i U_j)$ which implies that $k + 1 \in \mathcal{L}(A)$, a contradiction. If $|\text{gcd}(U_i, U_j)| = 1$, then $U_i U_j = W_1 W_2$ with $W_1, W_2 \in \mathcal{A}(G)$, $|W_1| = 2$, and $|W_2| = 4$, a contradiction to the maximality condition in Equation (14). Therefore $|\text{gcd}(U_i, U_j)| = 0$. This completes the proof of A. \square

Note that A.5 implies that $\{|U_i| \mid i \in [1, k_0]\} \neq \{3, 4, 5\}$. Thus it remains to discuss the following six subcases.

CASE 4.1. $\{|U_i| \mid i \in [1, k_0]\} = \{3, 5\}$.

By A.5 and A.7, we obtain that $|U_1| = 5, |U_2| = \dots = |U_{k_0}| = 3$, and that $U_1 \cdot \dots \cdot U_{k_0}$ is square-free. This implies that $\max \mathcal{L}(U_1 \cdot \dots \cdot U_{k_0}) = k_0$, and hence $\max \mathcal{L}(A) = \max \mathcal{L}(U_0 \cdot \dots \cdot U_{k_0}) + k - k_0 = k$, a contradiction.

CASE 4.2. $\{|U_i| \mid i \in [1, k_0]\} = \{3, 4\}$.

By A.6 and A.7, we obtain that $|U_1| = 4, |U_2| = \dots = |U_{k_0}| = 3$, and that $U_1 \cdot \dots \cdot U_{k_0}$ is square-free. This implies that $\max \mathcal{L}(U_1 \cdot \dots \cdot U_{k_0}) = k_0$, and hence $\max \mathcal{L}(A) = \max \mathcal{L}(U_0 \cdot \dots \cdot U_{k_0}) + k - k_0 = k$, a contradiction.

CASE 4.3. $\{|U_i| \mid i \in [1, k_0]\} = \{3\}$.

By A.7, we obtain that $U_1 \cdot \dots \cdot U_{k_0}$ is square-free. This implies that $\max \mathcal{L}(U_1 \cdot \dots \cdot U_{k_0}) = k_0$, and hence $\max \mathcal{L}(A) = \max \mathcal{L}(U_0 \cdot \dots \cdot U_{k_0}) + k - k_0 = k$, a contradiction.

CASE 4.4. $\{|U_i| \mid i \in [1, k_0]\} = \{5\}$.

By A.2, it follows that $A = U_1^{k_0} U_{k_0+1} \cdot \dots \cdot U_k$. If $\text{supp}(U_{k_0+1} \cdot \dots \cdot U_k) \subset \text{supp}(U_1)$, then $\Delta(\mathcal{L}(A)) = \{3\}$, a contradiction. Thus there exists $j \in [k_0 + 1, k]$ such that $U_j = g^2$ for some $g \notin \text{supp}(U_1)$. Then there exist $g_1, g_2 \in G$ such that $g_1 g_2 \mid U_1$ and $g =$

$g_1 + g_2$. It follows that $U_1^2 U_j = g_1^2 g_2^2 (U_1(g_1 g_2)^{-1} g)^2$, where $g_1^2, g_2^2, U_1(g_1 g_2)^{-1} g$ are atoms. Therefore $4 \in \mathcal{L}(U_1^2 U_j)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction.

CASE 4.5. $\{|U_i| \mid i \in [1, k_0]\} = \{4\}$.

Assume to the contrary, that $k_0 \geq 3$. Then **A.4** implies that $U_1 \cdots U_{k_0} = U_1^{k_0}$, and we set $G_1 = \langle \text{supp}(U_1) \rangle$. If there exists $g \in \text{supp}(U_{k_0+1} \cdots U_k)$ such that $g \in G_1 \setminus \text{supp}(U_1)$, then $4 \in \mathcal{L}(U_1^2 g^2)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. If there exist distinct $g_1, g_2 \in \text{supp}(U_{k_0+1} \cdots U_k)$ such that $g_1 \notin G_1$ and $g_2 \notin G_1$, then $g_1 + g_2 \in G_1$. Since $g_1 + g_2 \in \text{supp}(U_1)$ implies that $5 \in \mathcal{L}(U_1^2 g_1^2 g_2^2)$ and $k+1 \in \mathcal{L}(A)$, we obtain that $g_1 + g_2 \in G_1 \setminus \text{supp}(U_1)$. Then $U_1^2 g_1^2 g_2^2 = W_1^2 W_2 W_3$ where $W_1, W_2, W_3 \in \mathcal{A}(G)$ with $|W_1| = 4$, $W_1 \neq U_1$, and $|W_2| = |W_3| = 2$. Say $U_{k_0+1} = g_1^2$ and $U_{k_0+2} = g_2^2$. Then $W_1^2 U_3 \cdots U_{k_0} W_1 W_2 U_{k_0+3} \cdots U_k$ is a factorization of A of length k satisfying the maximality condition of Equation (14) and hence applying **A.4** to this factorization, we obtain a contradiction. Therefore $\text{supp}(U_{k_0+1} \cdots U_k) \subset \text{supp}(U_1) \cup \{g\}$ where g is independent from $\text{supp}(U_1)$ and hence $\text{supp}(A) \subset \text{supp}(U_1) \cup \{g\}$ which implies that $\Delta(\mathcal{L}(A)) = \{2\}$, a contradiction.

Therefore it follows that $k_0 = 2$. Then $U_1 = U_2$ (since otherwise we would have $\max \mathcal{L}(A) = k$ by $U_1 U_2$ is square-free), and we obtain that $\mathcal{L}(A) = [\min \mathcal{L}(A), k] \cup \{k+2\}$. Assume to the contrary that there exists a $W \in \mathcal{A}(G)$ such that $W \mid A$ and $|W| = 5$. Then there exist $g, g_1, g_2 \in G$ such that $g \mid U_1$, $g_1 g_2 \mid W$, and $g = g_1 + g_2$, and hence $|\{g_1, g_2\} \cap \text{supp}(U_1)| \leq 1$. If $\{g_1, g_2\} \cap \text{supp}(U_1) = \emptyset$, then there exist distinct $t_1, t_2 \in [k_0+1, k]$ such that $U_{t_1} = g_1^2$ and $U_{t_2} = g_2^2$. Thus $5 \in \mathcal{L}(U_1 U_2 U_{t_1} U_{t_2})$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction. Suppose that $|\{g_1, g_2\} \cap \text{supp}(U_1)| = 1$, say $g_1 \notin \text{supp}(U_1)$ and $g_2 \in \text{supp}(U_1)$. Then there exists $t \in [k_0+1, k]$ such that $U_t = g_1^2$. Therefore $4 \in \mathcal{L}(U_1 U_2 U_t)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction.

Thus every atom W with $W \mid A$ has length $|W| < 5$. It follows that $\min \mathcal{L}(A) \geq \lceil \frac{2 \max \mathcal{L}(A)}{4} \rceil = \lceil \frac{\max \mathcal{L}(A)}{2} \rceil$ and hence $\mathcal{L}(A) \in \mathcal{L}_5$.

CASE 4.6. $\{|U_i| \mid i \in [1, k_0]\} = \{4, 5\}$.

By **A.2**, **A.3**, and **A.4**, we obtain that $|\{U_1, \dots, U_{k_0}\}| = 2$. Without restriction we may assume that $U_1 \cdots U_{k_0} = U^{k_1} V^{k_2}$ where $k_1, k_2 \in \mathbb{N}$ with $k_0 = k_1 + k_2$ and $V = e_1 e_2 e_3 (e_1 + e_2 + e_3)$ (recall that (e_1, \dots, e_4) is a basis of G , $e_0 = e_1 + e_2 + e_3 + e_4$, and $U = e_1 e_2 e_3 e_4 e_0$). We claim that

- $\text{supp}(U_{k_0+1} \cdots U_k) \subset \text{supp}(UV)$.
- If $k_1 \geq 2$, then $\text{supp}(U_{k_0+1} \cdots U_k) \subset \text{supp}(U)$, and
- if $k_2 \geq 2$, then $\{e_4, e_0\} \not\subset \text{supp}(U_{k_0+1} \cdots U_k)$.

Indeed, assume to the contrary that $g \in \text{supp}(U_{k_0+1} \cdots U_k) \setminus \text{supp}(UV)$. By symmetry, we only need to consider $g = e_1 + e_2$ and $g = e_1 + e_4$ and both cases imply that $4 \in \mathcal{L}(UVg^2)$, a contradiction to $k+1 \notin \mathcal{L}(A)$. If $k_1 \geq 2$ and $g = e_1 + e_2 + e_3 \in \text{supp}(U_{k_0+1} \cdots U_k)$, then $4 \in \mathcal{L}(U^2 g^2)$ and $k+1 \in \mathcal{L}(A)$, a contradiction. Thus if $k_1 \geq 2$, then $\text{supp}(U_{k_0+1} \cdots U_k) \subset \text{supp}(U)$. If $k_2 \geq 2$ and $\{e_4, e_0\} \subset \text{supp}(U_{k_0+1} \cdots U_k)$, then $5 \in \mathcal{L}(V^2 e_4^2 e_0^2)$ and hence $k+1 \in \mathcal{L}(A)$, a contradiction.

Thus all three claims are proved, and we distinguish three subcases.

CASE 4.6.1. $k_1 = 1$.

If $\{e_4, e_0\} \not\subset \text{supp}(U_{k_0+1} \cdots U_k)$, then $L(A) = L(UV^{k_2}) + k - k_0 = L(V^{k_0}) + k - k_0$ and hence $\Delta(L(A)) = \{2\}$, a contradiction. If $\{e_4, e_0\} \subset \text{supp}(U_{k_0+1} \cdots U_k)$, then $k_2 = 1$ and we may assume that $U_{k_0+1} = e_4^2$ and that $U_{k_0+2} = e_0^2$. Then $L(A) = L(UVU_{k_0+1}U_{k_0+2}) + k - k_0 - 2 = \{k-1, k, k+2\}$ with $k \geq 4$, and hence $L(A) \in \mathcal{L}_5$.

CASE 4.6.2. $k_1 \geq 2$ and $k_2 \geq 2$.

Thus $\text{supp}(U_{k_0+1} \cdots U_k)$ is independent and it follows that $\text{supp}(U_{k_0+1} \cdots U_k) \subset \{e_1, e_2, e_3, e_4\}$ or $\text{supp}(U_{k_0+1} \cdots U_k) \subset \{e_1, e_2, e_3, e_0\}$. Then we have $L(A) = L(U^{k_1}V^{k_2}) + k - k_0$. By Equation (11), $L(U^{k_1}V^{k_2})$ is equal to

$$\begin{cases} \{k_0\} \cup [k_0 + 2, 5\lfloor k_1/2 \rfloor + 4(k_0/2 - \lfloor k_1/2 \rfloor)] & \text{if } k_0 = k_1 + k_2 \text{ is even,} \\ \{k_0\} \cup [k_0 + 2, 5\lfloor k_1/2 \rfloor + 4((k_0 - 1)/2 - \lfloor k_1/2 \rfloor) + 1] & \text{if } k_0 = k_1 + k_2 \text{ is odd.} \end{cases}$$

Let $\ell = \max L(U^{k_1}V^{k_2}) - k_0 - 2$ and hence

$$\ell = \begin{cases} k_0 + \lfloor \frac{k_1}{2} \rfloor - 2 & \text{if } k_0 \geq 4 \text{ is even,} \\ k_0 + \lfloor \frac{k_1}{2} \rfloor - 3 & \text{if } k_0 \geq 5 \text{ is odd.} \end{cases}$$

Since $k_1 \geq 2$ and $k_2 \geq 2$, we obtain that $\ell \geq 3$ and $\ell \neq 4$. We also have that

$$\ell \leq \begin{cases} k_0 + \lfloor \frac{k_0 - 2}{2} \rfloor - 2 = \frac{3k_0}{2} - 3 & \text{if } k_0 \text{ is even,} \\ k_0 + \lfloor \frac{k_0 - 2}{2} \rfloor - 3 = \frac{3k_0 - 9}{2} & \text{if } k_0 \text{ is odd.} \end{cases}$$

Therefore

$$k_0 \geq \begin{cases} \frac{2\ell}{3} + 2 & \text{if } k_0 \text{ is even,} \\ \frac{2\ell}{3} + 3 & \text{if } k_0 \text{ is odd,} \end{cases}$$

and hence

$$k_0 \geq \begin{cases} 2\lceil \frac{\ell}{3} \rceil + 2 & \text{if } k_0 \text{ is even,} \\ 2\lceil \frac{\ell}{3} \rceil + 2 & \text{if } k_0 \text{ is odd.} \end{cases}$$

It follows that $L(U^{k_1}V^{k_2}) \in \mathcal{L}_6$ which implies that $L(A) \in \mathcal{L}_6$.

CASE 4.6.3. $k_1 \geq 2$ and $k_2 = 1$.

Then $\text{supp}(U_{k_0+1} \cdots U_k) \subset \{e_1, e_2, e_3, e_4, e_0\}$. If $\{e_4, e_0\} \not\subset \text{supp}(U_{k_0+1} \cdots U_k)$, then $L(A) = L(U^{k_1}V) + k - k_0$ is equal to

$$\begin{cases} k + \{0, 2, 3\} + 3 \cdot [0, k_1/2 - 1], & \text{if } k_1 \text{ is even,} \\ k + \{0, 2, 3\} + 3 \cdot [0, (k_1 - 1)/2 - 1] \cup \{k + (3k_1 - 3)/2 + 2\}, & \text{if } k_1 \text{ is odd} \end{cases}$$

by Equation (12). Therefore $\mathsf{L}(A) \in \mathcal{L}_8$.

If $\{e_4, e_0\} \subset \text{supp}(U_{k_0+1} \cdot \dots \cdot U_k)$, then we may assume that $U_{k_0+1} = e_4^2$ and that $U_{k_0+2} = e_0^2$. Thus

$$\begin{aligned}\mathsf{L}(A) &= \mathsf{L}(U^{k_1} V U_{k_0+1} U_{k_0+2}) + k - k_0 - 2 \\ &= \begin{cases} k - 1 + \{0, 1, 3\} + 3 \cdot [0, (k_1 + 1)/2 - 1], & \text{if } k_1 \text{ is odd,} \\ k - 1 + \{0, 1, 3\} + 3 \cdot [0, k_1/2 - 1] \cup \{k + 3k_1/2 + 1\}, & \text{if } k_1 \text{ is even,} \end{cases}\end{aligned}$$

by Equation (13) and hence $\mathsf{L}(A) \in \mathcal{L}_7$. \square

5 Sets of lengths of weakly Krull monoids

It is well-known that – under reasonable algebraic finiteness conditions – the Structure Theorem for Sets of Lengths holds for weakly Krull monoids (as it is true for transfer Krull monoids of finite type, see Proposition 3.2). In spite of this common feature we will demonstrate that systems of sets of lengths for a variety of classes of weakly Krull monoids are different from the system of sets of lengths of any transfer Krull monoid (apart from well-described exceptional cases; see Theorems 5.5 to 5.8). Since half-factorial monoids are transfer Krull monoids, and since there are half-factorial weakly Krull monoids, half-factoriality is such a natural exceptional case.

So far there are only a couple of results in this direction. In [15], Frisch showed that $\text{Int}(\mathbb{Z})$, the ring of integer-valued polynomials over \mathbb{Z} , is not a transfer Krull domain (nevertheless, the system of sets of lengths of $\text{Int}(\mathbb{Z})^\bullet$ coincides with $\mathcal{L}(G)$ for an infinite abelian group G). To mention a result by Smertnig, let \mathcal{O} be the ring of integers of an algebraic number field K , A a central simple algebra over K , and R a classical maximal \mathcal{O} -order of A . Then R is a non-commutative Dedekind domain and in particular an HNP ring (see [32, Sections 5.2 and 5.3]). Furthermore, R is a transfer Krull domain if and only if every stably free left R -ideal is free ([35, Theorems 1.1 and 1.2]).

We gather basic concepts and properties of weakly Krull monoids and domains (Propositions 5.1 and 5.2). In the remainder of this section, all monoids and domains are supposed to be commutative.

Let H be a monoid (hence commutative, cancellative, and with unit element). We denote by $\mathsf{q}(H)$ the quotient group of H , by $H_{\text{red}} = H/H^\times$ the associated reduced monoid of H , by $\mathfrak{X}(H)$ the set of minimal nonempty prime s -ideals of H , and by $\mathfrak{m} = H \setminus H^\times$ the maximal s -ideal. Let $\mathcal{I}_v^*(H)$ denote the monoid of v -invertible v -ideals of H (with v -multiplication). Then $\mathcal{F}_v(H)^\times = \mathsf{q}(\mathcal{I}_v^*(H))$ is the quotient group of fractional v -invertible v -ideals, and $\mathcal{C}_v(H) = \mathcal{F}_v(H)^\times / \{xH \mid x \in \mathsf{q}(H)\}$ is the v -class group of H (detailed presentations of ideal theory in commutative monoids can be found in [30, 20]). We denote by $\hat{H} \subset \mathsf{q}(H)$ the complete integral

closure of H , and by $(H : \widehat{H}) = \{x \in \mathbf{q}(H) \mid x\widehat{H} \subset H\} \subset H$ the conductor of H . A submonoid $S \subset H$ is said to be saturated if $S = \mathbf{q}(S) \cap H$. For the definition and discussion of the concepts of being faithfully saturated or being locally tame we refer to [20, Sections 1.6 and 3.6].

To start with the local case, we recall that H is said to be

- *primary* if $\mathbf{m} \neq \emptyset$ and for all $a, b \in \mathbf{m}$ there is an $n \in \mathbb{N}$ such that $b^n \subset aH$.
- *strongly primary* if $\mathbf{m} \neq \emptyset$ and for every $a \in \mathbf{m}$ there is an $n \in \mathbb{N}$ such that $\mathbf{m}^n \subset aH$. We denote by $\mathcal{M}(a)$ the smallest n having this property.
- a *discrete valuation monoid* if it is primary and contains a prime element (equivalently, $H_{\text{red}} \cong (\mathbb{N}_0, +)$).

Furthermore, H is said to be

- *weakly Krull* ([30, Corollary 22.5]) if

$$H = \bigcap_{\mathbf{p} \in \mathfrak{X}(H)} H_{\mathbf{p}} \quad \text{and} \quad \{\mathbf{p} \in \mathfrak{X}(H) \mid a \in \mathbf{p}\} \quad \text{is finite for all } a \in H.$$

- *weakly factorial* if one of the following equivalent conditions is satisfied ([30, Exercise 22.5]):

- Every non-unit is a finite product of primary elements.
- H is a weakly Krull monoid with trivial t -class group.

Clearly, every localization $H_{\mathbf{p}}$ of H at a minimal prime ideal $\mathbf{p} \in \mathfrak{X}(H)$ is primary, and a weakly Krull monoid H is v -noetherian if and only if $H_{\mathbf{p}}$ is v -noetherian for each $\mathbf{p} \in \mathfrak{X}(H)$. Every v -noetherian primary monoid H is strongly primary and v -local, and if $(H : \widehat{H}) \neq \emptyset$, then H is locally tame ([21, Lemma 3.1 and Corollary 3.6]). Every strongly primary monoid is a primary BF-monoid ([20, Section 2.7]). Therefore the coproduct of a family of strongly primary monoids is a BF-monoid, and every coproduct of a family of primary monoids is weakly factorial. A v -noetherian weakly Krull monoid H is weakly factorial if and only if $\mathcal{C}_v(H) = 0$ if and only if $H_{\text{red}} \cong \mathcal{J}_v^*(H)$.

By a numerical monoid H we mean an additive submonoid of $(\mathbb{N}_0, +)$ such that $\mathbb{N}_0 \setminus H$ is finite. Clearly, every numerical monoid is v -noetherian primary, and hence it is strongly primary. Note that a numerical monoid is half-factorial if and only if it is equal to $(\mathbb{N}_0, +)$.

Let R be a domain. Then $R^\bullet = R \setminus \{0\}$ is a monoid, and all arithmetic and ideal theoretic concepts introduced for monoids will be used for domains in the obvious way. The domain R is weakly Krull (resp. weakly factorial) if and only if its multiplicative monoid R^\bullet is weakly Krull (resp. weakly factorial). Weakly Krull domains were introduced by Anderson, Anderson, Mott, and Zafrullah ([2, 3]). We recall some most basic facts and refer to an extended list of weakly Krull domains and monoids in [22, Examples 5.7]. The monoid R^\bullet is primary if and only if R is one-dimensional and local. If R is one-dimensional local Mori, then R^\bullet is strongly primary and locally tame ([23]). Furthermore, every one-dimensional semilocal Mori domain with nontrivial conductor is weakly factorial and the same holds true for

generalized Cohen-Kaplansky domains. It can be seen from the definition that one-dimensional noetherian domains are v -noetherian weakly Krull domains.

Proposition 5.1 summarizes the main algebraic properties of v -noetherian weakly Krull monoids and Proposition 5.2 recalls that their arithmetic can be studied via weak transfer homomorphisms to weakly Krull monoids of very special form.

Proposition 5.1. *Let H be a v -noetherian weakly Krull monoid.*

1. *The monoid $\mathcal{I}_v^*(H)$ is isomorphic to the coproduct of $(H_{\mathfrak{p}})_{\text{red}}$ over all $\mathfrak{p} \in \mathfrak{X}(H)$. In particular, $\mathcal{I}_v^*(H)$ is weakly factorial and v -noetherian.*
2. *Suppose that $\mathfrak{f} = (H : \widehat{H}) \neq \emptyset$. We set $\mathcal{P}^* = \{\mathfrak{p} \in \mathfrak{X}(H) \mid \mathfrak{p} \supset \mathfrak{f}\}$, and $\mathcal{P} = \mathfrak{X}(H) \setminus \mathcal{P}^*$.*
 - a. *Then \widehat{H} is Krull, \mathcal{P}^* is finite, and $H_{\mathfrak{p}}$ is a discrete valuation monoid for each $\mathfrak{p} \in \mathcal{P}$. In particular, $\mathcal{I}_v^*(H)$ is isomorphic to $\mathcal{F}(\mathcal{P}) \times \prod_{\mathfrak{p} \in \mathcal{P}^*} (H_{\mathfrak{p}})_{\text{red}}$.*
 - b. *If $\mathcal{H} = \{aH \mid a \in H\}$ is the monoid of principal ideals of H , then $\mathcal{H} \subset \mathcal{I}_v^*(H)$ is saturated. Moreover, if H is the multiplicative monoid of a domain, then all monoids $H_{\mathfrak{p}}$ are locally tame and $\mathcal{H} \subset \mathcal{I}_v^*(H)$ is faithfully saturated.*

Proof. 1. See [22, Proposition 5.3].

2. For (a) we refer to [20, Theorem 2.6.5] and for (b) we refer to [20, Theorems 3.6.4 and 3.7.1]. \square

Proposition 5.2. *Let $D = \mathcal{F}(\mathcal{P}) \times \prod_{i=1}^n D_i$ be a monoid, where $\mathcal{P} \subset D$ is a set of primes, $n \in \mathbb{N}_0$, and D_1, \dots, D_n are reduced primary monoids. Let $H \subset D$ be a saturated submonoid, $G = \mathfrak{q}(D)/\mathfrak{q}(H)$, and $G_{\mathcal{P}} = \{p\mathfrak{q}(H) \mid p \in \mathcal{P}\} \subset G$ the set of classes containing primes.*

1. *There is a saturated submonoid $B \subset F = \mathcal{F}(G_{\mathcal{P}}) \times \prod_{i=1}^n D_i$ and a weak transfer homomorphism $\theta: H \rightarrow B$. Moreover, if G is a torsion group, then there is a monomorphism $\mathfrak{q}(F)/\mathfrak{q}(B) \rightarrow G$.*
2. *If G is a torsion group, then H is weakly Krull.*

Proof. 1. See [20, Propositions 3.4.7 and 3.4.8].

2. See [22, Lemma 5.2]. \square

Note that, under the assumption of 5.1.2, the embedding $\mathcal{H} \hookrightarrow \mathcal{I}_v^*(H)$ fulfills the assumptions imposed on the embedding $H \hookrightarrow D$ in Proposition 5.2. Thus Proposition 5.2 applies to v -noetherian weakly Krull monoids. For simplicity and in order to avoid repetitions, we formulate the next results (including Theorem 5.7) in the abstract setting of Proposition 5.2. However, v -noetherian weakly Krull domains and their monoids of v -invertible v -ideals are in the center of our interest.

If (in the setting of Proposition 5.2) $G_{\mathcal{P}}$ is finite, then $F = \mathcal{F}(G_{\mathcal{P}}) \times \prod_{i=1}^n D_i$ is a finite product of primary monoids and $B \subset F$ is a saturated submonoid. We formulate the main structural result for sets of lengths in v -noetherian weakly Krull monoids in this abstract setting.

Proposition 5.3. *Let D_1, \dots, D_n be locally tame strongly primary monoids and $H \subset D = D_1 \times \dots \times D_n$ a faithfully saturated submonoid such that $\mathbf{q}(D)/\mathbf{q}(H)$ is finite.*

1. *The monoid H satisfies the Structure Theorem for Sets of Lengths.*
2. *There is a finite abelian group G such that for every $L \in \mathcal{L}(H)$ there is a $y \in \mathbb{N}$ such that $y + L \in \mathcal{L}(G)$.*

Proof. 1. follows from [20, Theorem 4.5.4], and 2. follows from 1. and from Proposition 3.2.2. \square

The next lemma on zero-sum sequences will be crucial in order to distinguish between sets of lengths in weakly Krull monoids and sets of lengths in transfer Krull monoids.

Lemma 5.4. *Let G be an abelian group and $G_0 \subset G$ a non-half-factorial subset.*

1. *There exists a half-factorial subset $G_1 \subset G_0$ with $\mathcal{B}(G_1) \neq \{1\}$.*
2. *There are $M \in \mathbb{N}$ and zero-sum sequences $B_k \in \mathcal{B}(G_0)$ for every $k \in \mathbb{N}$ such that $2 \leq |\mathcal{L}(B_k)| \leq M$ but $\min \mathcal{L}(B_k) \rightarrow \infty$ as $k \rightarrow \infty$.*

Proof. 1. Since G_0 is not half-factorial, there is a $B \in \mathcal{B}(G_0)$ such that $|\mathcal{L}(B)| > 1$. Thus $\text{supp}(B)$ is finite and not half-factorial, say $\text{supp}(B) = \{g_1, \dots, g_\ell\}$ with $\ell \geq 2$. Without restriction we may suppose that every proper subset of $\{g_1, \dots, g_\ell\}$ is half-factorial. Assume to the contrary that for every subset $G_1 \subsetneq \{g_1, \dots, g_\ell\}$ we have $\mathcal{B}(G_1) = \{1\}$. Since $\{g_1, \dots, g_\ell\}$ is minimal non-half-factorial, there is an atom $A_1 \in \mathcal{A}(\{g_1, \dots, g_\ell\})$ such that $v_{g_i}(A_1) > 0$ for every $i \in [1, \ell]$. Since $\{g_1, \dots, g_\ell\}$ is not half-factorial, there is an atom $A_2 \in \mathcal{A}(\{g_1, \dots, g_\ell\})$ distinct from A_1 , say

$$A_1 = g_1^{k_1} \cdots g_\ell^{k_\ell} \quad \text{and} \quad A_2 = g_1^{t_1} \cdots g_\ell^{t_\ell}$$

where $k_i \in \mathbb{N}$ and $t_i \in \mathbb{N}_0$ for every $i \in [1, \ell]$. Let $\tau \in [1, \ell]$ such that $\frac{t_\tau}{k_\tau} = \max\{\frac{t_j}{k_j} \mid j \in [1, \ell]\}$. Then $k_j t_\tau - t_j k_\tau \geq 0$ for every $j \in [1, \ell]$ whence

$$W = A_2^{t_\tau} A_1^{-k_\tau} \in \mathcal{B}(\{g_1, \dots, g_\ell\} \setminus \{g_\tau\}),$$

which implies that $W = 1$. Therefore $\frac{t_\tau}{k_\tau} = \frac{t_j}{k_j}$ for every $j \in [1, \ell]$ and hence $A_1 \mid A_2$ or $A_2 \mid A_1$, a contradiction.

2. Let $B \in \mathcal{B}(G_0)$ with $|\mathcal{L}(B)| > 1$. By 1., there exists a half-factorial subset $G_1 \subsetneq G_0$ such that $\mathcal{B}(G_1) \neq \{1\}$. Let $A \in \mathcal{A}(G_1)$ and $B_k = A^k B$ for every $k \in \mathbb{N}$. Obviously there exists $k_0 \in \mathbb{N}$ such that $\mathcal{L}(B_k) = \mathcal{L}(A^{k-k_0}) + \mathcal{L}(B_{k_0}) = k - k_0 + \mathcal{L}(B_{k_0})$ for every $k \geq k_0$. Thus $|\mathcal{L}(B_k)| \leq \max \mathcal{L}(B_{k_0}) - \min \mathcal{L}(B_{k_0})$ and $\min \mathcal{L}(B_k) = k - k_0 + \min \mathcal{L}(B_{k_0})$. \square

Now we consider strongly primary monoids and work out a feature of their systems of sets of lengths which does not occur in the system of sets of lengths of any transfer Krull monoid. To do so we study the set $\{\rho(L) \mid L \in \mathcal{L}(H)\}$ of elasticities

of all sets of lengths. This set was studied first by Chapman et al. in a series of papers (see [6, 13, 7, 8]). Among others they showed that in an atomic monoid H , which has a prime element and an element $a \in H$ with $\rho(L(a)) = \rho(H)$, every rational number q with $1 \leq q \leq \rho(H)$ can be realized as the elasticity of some $L \in \mathcal{L}(H)$ ([6, Corollary 2.2]). Primary monoids, which are not discrete valuation monoids, have no prime elements and their set of elasticities is different, as we will see in the next theorem. Statement 1. of Theorem 5.5 was proved for numerical monoids in [13, Theorem 2.2].

Theorem 5.5. *Let H be a strongly primary monoid that is not half-factorial.*

1. *There is a $\beta \in \mathbb{Q}_{>1}$ such that $\rho(L) \geq \beta$ for all $L \in \mathcal{L}(H)$ with $\rho(L) \neq 1$.*
2. *$\mathcal{L}(H) \neq \mathcal{L}(G_0)$ for any subset G_0 of any abelian group. In particular, H is not a transfer Krull monoid.*
3. *If one of the following two conditions holds, then H is locally tame.*
 - $\sup\{\min L(c) \mid c \in H\} < \infty$.
 - *There exists some $u \in H \setminus H^\times$ such that $\rho_{\mathcal{M}(u)}(H) < \infty$.*

If H is locally tame, then $\Delta(H)$ is finite, and there is an $M \in \mathbb{N}_0$ such that every $L \in \mathcal{L}(H)$ is an AAMP with period $\{0, \min \Delta(H)\}$ and bound M .

Remark. If H is the multiplicative monoid of a one-dimensional local Mori domain R with nonzero conductor $(R:\widehat{R}) \neq \{0\}$, then one of the conditions in 3. is satisfied (see [20, Proposition 2.10.7 and Theorem 3.1.5]). However, there are strongly primary monoids for which none of the conditions holds and which are not locally tame ([21, Proposition 3.7]).

Proof. 1. Let $b \in H$ such that $|L(b)| \geq 2$ and let $u \in \mathcal{A}(H)$. Since H is a strongly primary monoid, we have $(H \setminus H^\times)^{\mathcal{M}(b)} \in bH$ and $(H \setminus H^\times)^{\mathcal{M}(u)} \in uH$. Thus $b \mid u^{\mathcal{M}(b)}$ and hence $|L(u^{\mathcal{M}(b)})| \geq 2$. We define

$$\beta_1 = \frac{\mathcal{M}(b) + \mathcal{M}(u) + 1}{\mathcal{M}(b) + \mathcal{M}(u)}, \quad \beta_2 = \frac{\max L(u^{\mathcal{M}(b)}) + \mathcal{M}(b) + \mathcal{M}(u)}{\min L(u^{\mathcal{M}(b)}) + \mathcal{M}(b) + \mathcal{M}(u)},$$

and observe that $\beta = \min\{\beta_1, \beta_2\} > 1$. Let $a \in H$ with $\rho(L(a)) \neq 1$. We show that $\rho(L(a)) \geq \beta$.

Let $k \in \mathbb{N}_0$ be maximal such that $u^k \mid a$, say $a = u^k u'$ with $u' \in H$. Thus $u \nmid u'$ and thus $\max L(u') < \mathcal{M}(u)$. If $k < \mathcal{M}(b)$, then $\min L(a) \leq \min L(u^k) + \min L(u') \leq \mathcal{M}(b) + \mathcal{M}(u)$, and hence

$$\rho(L(a)) = \frac{\max L(a)}{\min L(a)} \geq \frac{\min L(a) + 1}{\min L(a)} \geq \frac{\mathcal{M}(b) + \mathcal{M}(u) + 1}{\mathcal{M}(b) + \mathcal{M}(u)} = \beta_1 \geq \beta.$$

If $k \geq \mathcal{M}(b)$, then there exist $t \in \mathbb{N}$ and $t_0 \in [0, \mathcal{M}(b) - 1]$ such that $k = t\mathcal{M}(b) + t_0$, and hence

$$\begin{aligned}
\rho(\mathbf{L}(a)) &= \frac{\max \mathbf{L}(a)}{\min \mathbf{L}(a)} \geq \frac{\max \mathbf{L}(u^k) + \max \mathbf{L}(u')}{\min \mathbf{L}(u^k) + \min \mathbf{L}(u')} \\
&\geq \frac{t \max \mathbf{L}(u^{\mathcal{M}(b)}) + \max \mathbf{L}(u^{t_0}) + \max \mathbf{L}(u')}{t \min \mathbf{L}(u^{\mathcal{M}(b)}) + \min \mathbf{L}(u^{t_0}) + \min \mathbf{L}(u')} \\
&\geq \frac{t \max \mathbf{L}(u^{\mathcal{M}(b)}) + t_0 + \max \mathbf{L}(u')}{t \min \mathbf{L}(u^{\mathcal{M}(b)}) + t_0 + \max \mathbf{L}(u')} \\
&\geq \frac{t \max \mathbf{L}(u^{\mathcal{M}(b)}) + \mathcal{M}(b) + \mathcal{M}(u)}{t \min \mathbf{L}(u^{\mathcal{M}(b)}) + \mathcal{M}(b) + \mathcal{M}(u)} \geq \beta_2 \geq \beta.
\end{aligned}$$

2. Assume to the contrary that there are an abelian group G and a subset $G_0 \subset G$ such that $\mathcal{L}(H) = \mathcal{L}(G_0)$. Since H is not half-factorial, G_0 is not half-factorial. By 1., there exists $\beta \in \mathbb{Q}$ with $\beta > 1$ such that $\rho(L) \geq \beta$ for every $L \in \mathcal{L}(H)$. Lemma 5.4.2 implies that there are zero-sum sequences $B_k \in \mathcal{B}(G_0)$ such that $\rho(\mathbf{L}(B_k)) \rightarrow 1$ as $k \rightarrow \infty$, a contradiction.

3. This follows from [20, 3.1.1, 3.1.2, and 4.3.6]. \square

Sets of lengths of numerical monoids have found wide attention in the literature (see, among others, [9, 1, 14]). As can be seen from Theorem 5.5.3, the structure of their sets of lengths is simpler than the structure of sets of lengths of transfer Krull monoids over finite abelian groups. Thus it is no surprise that there are infinitely many non-isomorphic numerical monoids whose systems of sets of lengths coincide, and that an analog of Conjecture 3.4 for numerical monoids does not hold true ([1]). It is open whether for every $d \in \mathbb{N}$ and every $M \in \mathbb{N}_0$ there is a strongly primary monoid D such that every AAMP with period $\{0, d\}$ and bound M can (up to a shift) be realized as a set of lengths in D (this would be the analog to the realization theorem given in Proposition 3.2.2). However, for every finite set $L \subset \mathbb{N}_{\geq 2}$ there is a v -noetherian primary monoid D and an element $a \in D$ such that $L = \mathbf{L}(a)$ ([21, Theorem 4.2]).

By Theorem 3.6 and Proposition 3.2.3, we know that $\{k, k+1\} \in \mathcal{L}(G)$ for every $k \geq 2$ and every abelian group G with $|G| \geq 3$. Furthermore, Theorem 3.7 is in sharp contrast to Theorem 5.6.1.

Theorem 5.6. *Let $D = D_1 \times \dots \times D_n$ be the direct product of strongly primary monoids D_1, \dots, D_n , which are not half-factorial.*

1. *For every finite nonempty set $L \subset \mathbb{N}$, there is a $y_L \in \mathbb{N}_0$ such that $y + L \notin \mathcal{L}(D)$ for any $y \geq y_L$.*
2. *We have $\mathcal{L}(D) \neq \mathcal{L}(G_0)$ for any subset G_0 of any abelian group, and hence D is not a transfer Krull monoid. If D_1, \dots, D_n are locally tame, then D satisfies the Structure Theorem for Sets of Lengths.*

Proof. For every $i \in [1, n]$ we choose an element $a_i \in D_i$ such that $|\mathbf{L}(a_i)| > 1$.

1. Let $L \subset \mathbb{N}$ be a finite nonempty set and let $y_L = |L|(\mathcal{M}(a_1) + \dots + \mathcal{M}(a_n))$. Assume to the contrary that there are $y \geq y_L$ and an element $b = b_1 \cdot \dots \cdot b_n \in D$ such

that $L(b) = y + L$. Then there is an $i \in [1, n]$ such that $\min L(b_i) \geq |L|\mathcal{M}(a_i)$. Then $b_i \in (D_i \setminus D_i^\times)^{\min L(b_i)} \subset (D_i \setminus D_i^\times)^{|L|\mathcal{M}(a_i)} \subset a_i^{|L|} D_i$. Thus there is a $c_i \in D_i$ such that $a_i^{|L|} c_i = b_i$. This implies that $|L|L(a_i) + L(c_i) \subset L(b_i)$. Since $|L(a_i)| \geq 2$, we infer that $|L(b_i)| \geq |L| + 1$ and hence $|L| = |y + L| = |L(b)| \geq |L(b_i)| \geq |L| + 1$, a contradiction.

2. By 1. and Lemma 5.4.2, the first conclusion follows.

If D_1, \dots, D_n are locally tame, then D satisfies the Structure Theorem by Proposition 5.3.1. \square

Theorem 5.7. *Let $D = \mathcal{F}(\mathcal{P}) \times D_1$ be the direct product of a free abelian monoid with nonempty basis \mathcal{P} and of a locally tame strongly primary monoid D_1 , and let G be an abelian group. Then D satisfies the Structure Theorem for Sets of Lengths, and the following statements are equivalent:*

- (a) $\mathcal{L}(D) = \mathcal{L}(G)$.
- (b) One of the following cases holds :
 - (b1) $|G| \leq 2$ and $\rho(D) = 1$.
 - (b2) G is isomorphic either to C_3 or to $C_2 \oplus C_2$, $[2, 3] \in \mathcal{L}(D)$, $\rho(D) = 3/2$, and $\Delta(D) = \{1\}$.
 - (b3) G is isomorphic to $C_3 \oplus C_3$, $[2, 5] \in \mathcal{L}(D)$, $\rho(D) = 5/2$, and $\Delta(D) = \{1\}$.

Remark. Let H be a v -noetherian weakly Krull monoid. If the conductor $(H : \widehat{H}) \in v\text{-max}(H)$, then by Proposition 5.1, $\mathcal{I}_v^*(H)$ is isomorphic to a monoid D as given in Theorem 5.7.

Proof. Since \mathcal{P} is nonempty, $\mathcal{L}(D) = \{y + L \mid y \in \mathbb{N}_0, L \in \mathcal{L}(D_1)\}$ whence $\Delta(D) = \Delta(D_1)$ and $\rho(D) = \rho(D_1)$. In particular, D is half-factorial if and only if D_1 is half-factorial. Since D_1 satisfies the Structure Theorem of Sets of Lengths by Theorem 5.5.3, the same is true for D .

If D is half-factorial and $\mathcal{L}(D) = \mathcal{L}(G)$, then $\rho(D) = \rho(D_1) = 1$ and G is half-factorial whence $|G| \leq 2$ by Proposition 3.3. Conversely, if $|G| \leq 2$ and $\rho(D) = 1$, then G and D are half-factorial and $\mathcal{L}(G) = \mathcal{L}(D)$.

Thus from now on we suppose that D_1 is not half-factorial and that (b1) does not hold. Then $\Delta(D) \neq \emptyset$ and we set $\min \Delta(D) = d$.

(a) \Rightarrow (b) Theorem 5.5.3 and Proposition 3.2.3 imply that G is finite. Since G is not half-factorial, it follows that $|G| \geq 3$. Theorem 5.5.3 shows that $\Delta_1(D) = \{d\}$, and since $1 \in \Delta_1(G) = \Delta_1(D)$, we infer that $d = 1$. Corollary 4.3.16 in [20] and [29, Theorem 1.1] imply that

$$\max\{\exp(G) - 2, r(G) - 1\} = \max \Delta_1(G) = \max \Delta_1(D) = 1.$$

Therefore G is isomorphic to one of the following groups: $C_2 \oplus C_2$, C_3 , $C_3 \oplus C_3$. We distinguish two cases.

CASE 1: G is isomorphic to $C_2 \oplus C_2$ or to C_3 .

By Proposition 3.3, we have

$$\mathcal{L}(D) = \mathcal{L}(C_2 \oplus C_2) = \mathcal{L}(C_3) = \{y + 2k + [0, k] \mid y, k \in \mathbb{N}_0\}.$$

In particular, we have $3/2 = \rho(G) = \rho(D)$ and $\{1\} = \Delta(G) = \Delta(D)$.

CASE 2: G is isomorphic to $C_3 \oplus C_3$.

By Theorem 4.1, just using different notation, we have

$$\begin{aligned} \mathcal{L}(D) = \mathcal{L}(C_3^2) &= \{[2k, \ell] \mid k \in \mathbb{N}_0, \ell \in [2k, 5k]\} \\ &\cup \{[2k+1, \ell] \mid k \in \mathbb{N}, \ell \in [2k+1, 5k+2]\} \cup \{\{1\}\}. \end{aligned}$$

In particular, we have $5/2 = \rho(G) = \rho(D)$ and $\{1\} = \Delta(G) = \Delta(D)$.

(b) \Rightarrow (a) First suppose that Case (b2) holds. We show that

$$\mathcal{L}(D) = \{y + 2k + [0, k] \mid y, k \in \mathbb{N}_0\}.$$

Then $\mathcal{L}(D) = \mathcal{L}(G)$ by Proposition 3.3. Since $\rho(D) = 3/2$ and $\Delta(D) = \{1\}$, it follows that $\mathcal{L}(D)$ is contained in the above family of sets. Thus we have to verify that for every $y, k \in \mathbb{N}_0$, the set $y + [2k, 3k] \in \mathcal{L}(D)$. Since \mathcal{P} is nonempty, D contains a prime element and hence it suffices to show that $[2k, 3k] \in \mathcal{L}(D)$ for all $k \in \mathbb{N}$. Let $a \in D$ with $L(a) = \{2, 3\}$, and let $k \in \mathbb{N}$. Then $\min L(a^k) \leq 2k$ and $\max L(a^k) \geq 3k$. Since $\rho(L(a^k)) \leq \rho(D) = 3/2$, it follows that $\min L(a^k) = 2k$ and $\max L(a^k) = 3k$. Since $\Delta(D) = \{1\}$, we finally obtain that $L(a^k) = [2k, 3k]$.

Now suppose that Case (b3) holds. We show that $\mathcal{L}(D)$ is equal to

$$\{[2k, \ell] \mid k \in \mathbb{N}_0, \ell \in [2k, 5k]\} \cup \{[2k+1, \ell] \mid k \in \mathbb{N}, \ell \in [2k+1, 5k+2]\} \cup \{\{1\}\}.$$

Then $\mathcal{L}(D) = \mathcal{L}(G)$ by Theorem 4.1. Since $\rho(D) = 5/2$ and $\Delta(D) = \{1\}$, it follows that $\mathcal{L}(D)$ is contained in the above family of sets. Now the proof runs along the same lines as the proof in Case (b2). \square

We show that the Cases (b2) and (b3) in Theorem 5.7 can actually occur. Recall that numerical monoids are locally tame and strongly primary. Let D_1 be a numerical monoid distinct from $(\mathbb{N}_0, +)$, say $\mathcal{A}(D_1) = \{n_1, \dots, n_t\}$ where $t \in \mathbb{N}_{\geq 2}$ and $1 < n_1 < \dots < n_t$. Then, by [13, Theorem 2.1] and [9, Proposition 2.9],

$$\rho(D_1) = \frac{n_t}{n_1} \quad \text{and} \quad \min \Delta(D_1) = \gcd(n_2 - n_1, \dots, n_t - n_{t-1}).$$

Suppose that $\rho(D_1) = m/2$ with $m \in \{3, 5\}$ and $\Delta(D_1) = \{1\}$. Then there is an $a \in D_1$ with $L(a) = [2, m] \in \mathcal{L}(D_1)$. Clearly, there are non-isomorphic numerical monoids with elasticity $m/2$ and set of distances equal to $\{1\}$.

Theorem 5.8. *Let R be a v-noetherian weakly Krull domain with conductor $\{0\} \subsetneq \mathfrak{f} = (R : \widehat{R}) \subsetneq R$, and let $\pi: \mathfrak{X}(\widehat{R}) \rightarrow \mathfrak{X}(R)$ be the natural map defined by $\pi(\mathfrak{P}) = \mathfrak{P} \cap R$ for all $\mathfrak{P} \in \mathfrak{X}(\widehat{R})$.*

1. a. $\mathcal{I}_v^*(H)$ is locally tame with finite set of distances, and it satisfies the Structure Theorem for Sets of Lengths.
- b. If π is not bijective, then $\mathcal{L}(\mathcal{I}_v^*(H)) \neq \mathcal{L}(G_0)$ for any finite subset G_0 of any abelian group and for any subset G_0 of an infinite cyclic group. In particular, $\mathcal{I}_v^*(H)$ is not a transfer Krull monoid of finite type.
- c. If R is seminormal, then the following statements are equivalent:
 - i. π is bijective.
 - ii. $\mathcal{I}_v^*(H)$ is a transfer Krull monoid of finite type.
 - iii. $\mathcal{I}_v^*(H)$ is half-factorial.

2. Suppose that the class group $\mathcal{C}_v(R)$ is finite.
 - a. The monoid R^\bullet of nonzero elements of R is locally tame with finite set of distances, and it satisfies the Structure Theorem for Sets of Lengths.
 - b. If π is not bijective, then $\mathcal{L}(R^\bullet) \neq \mathcal{L}(G_0)$ for any finite subset G_0 of any abelian group and for any subset G_0 of an infinite cyclic group. In particular, R is not a transfer Krull domain of finite type.
 - c. If π is bijective, R is seminormal, every class of $\mathcal{C}_v(R)$ contains a $\mathfrak{p} \in \mathfrak{X}(R)$ with $\mathfrak{p} \not\supset \mathfrak{f}$, and the natural epimorphism $\delta: \mathcal{C}_v(R) \rightarrow \mathcal{C}_v(\hat{R})$ is an isomorphism, then there is a weak transfer homomorphism $\theta: R^\bullet \rightarrow \mathcal{B}(\mathcal{C}_v(R))$. In particular, R is a transfer Krull domain of finite type.

Proof. Since $\mathfrak{f} \neq R$, it follows that $R \neq \hat{R}$ and that R is not a Krull domain. We use the structural description of $\mathcal{I}_v^*(H)$ as given in Proposition 5.1.

1.(a) and 2.(a) Both monoids, R^\bullet and $\mathcal{I}_v^*(H)$, are locally tame with finite set of distances by [20, Theorem 3.7.1]. Furthermore, they both satisfy the Structure Theorem for Sets of Lengths by Proposition 5.3 (use Propositions 5.1 and 5.2).

1.(b) and 2.(b) Suppose that π is not bijective. Then $\rho(\mathcal{I}_v^*(H)) = \rho(R^\bullet) = \infty$ by [20, Theorems 3.1.5 and 3.7.1]. Let G_0 be a finite subset of an abelian group G . Then $\mathcal{B}(G_0)$ is finitely generated, the Davenport constant $D(G_0)$ is finite whence the set of distances $\Delta(G_0)$ and the elasticity $\rho(G_0)$ are both finite (see [20, Theorems 3.4.2 and 3.4.11]). Thus $\mathcal{L}(\mathcal{I}_v^*(H)) \neq \mathcal{L}(G_0)$ and $\mathcal{L}(R^\bullet) \neq \mathcal{L}(G_0)$. If G_0 is a subset of an infinite cyclic group, then the set of distances is finite if and only if the elasticity is finite by [18, Theorem 4.2], and hence the assertion follows again.

1.(c) Suppose that R is seminormal. By 1.(b) and since half-factorial monoids are transfer Krull monoids of finite type, it remains to show that π is bijective if and only if $\mathcal{I}_v^*(H)$ is half-factorial. Since R is seminormal, all localizations $R_{\mathfrak{p}}$ with $\mathfrak{p} \in \mathfrak{X}(H)$ are seminormal. Thus $\mathcal{I}_v^*(H)$ is isomorphic to a monoid of the form $\mathcal{F}(\mathcal{P}) \times D_1 \times \dots \times D_n$, where $n \in \mathbb{N}$ and D_1, \dots, D_n are seminormal finitely primary monoids, and this monoid is half-factorial if and only if each monoid D_1, \dots, D_n is half-factorial. By [22, Lemma 3.6], D_i is half-factorial if and only if it has rank one for each $i \in [1, n]$, and this is equivalent to π being bijective (see [20, Theorem 3.7.1]).

2.(c) This follows from [22, Theorem 5.8]. \square

Note that every order R in an algebraic number field is a v -noetherian weakly Krull domain with finite class group $\mathcal{C}_v(R)$ such that every class contains a $\mathfrak{p} \in \mathfrak{X}(R)$ with $\mathfrak{p} \not\supseteq \mathfrak{f}$. If R is a v -noetherian weakly Krull domain as above, then Theorems 5.5, 5.6, and 5.7 provide further instances of when R is not a transfer Krull domain, but a characterization of the general case remains open. We formulate the following problem (see also [17, Problem 4.7]).

Problem 5.9. Let H be a v -noetherian weakly Krull monoid with nonempty conductor $(H:\widehat{H})$ and finite class group $\mathcal{C}_v(H)$. Characterize when H and when the monoid $\mathcal{J}_v^*(H)$ are transfer Krull monoids resp. transfer Krull monoids of finite type.

References

1. J. Amos, S.T. Chapman, N. Hine, and J. Paixão, *Sets of lengths do not characterize numerical monoids*, *Integers* **7** (2007), Paper A50, 8p.
2. D.D. Anderson, D.F. Anderson, and M. Zafrullah, *Atomic domains in which almost all atoms are prime*, *Commun. Algebra* **20** (1992), 1447 – 1462.
3. D.D. Anderson, J. Mott, and M. Zafrullah, *Finite character representations for integral domains*, *Boll. Unione Mat. Ital.* **6** (1992), 613 – 630.
4. D. Bachman, N. Baeth, and A. McQueen, *Factorizations of upper triangular Toeplitz matrices*, *Boll. Unione Mat. Ital.* **8** (2010), 131 – 150.
5. N.R. Baeth and D. Smertnig, *Factorization theory: From commutative to noncommutative settings*, *J. Algebra* **441** (2015), 475 – 551.
6. P. Baginski, S.T. Chapman, C. Crutchfield, K.G. Kennedy, and M. Wright, *Elastic properties and prime elements*, *Result. Math.* **49** (2006), 187 – 200.
7. P. Baginski, S.T. Chapman, M. Holden, and T. Moore, *Asymptotic elasticity in atomic monoids*, *Semigroup Forum* **72** (2006), 134 – 142.
8. T. Barron, C.O’Neill, and R. Pelayo, *On the set of elasticities in numerical monoids*, *Semigroup Forum* **94** (2017), 3750.
9. C. Bowles, S.T. Chapman, N. Kaplan, and D. Reiser, *On delta sets of numerical monoids*, *J. Algebra Appl.* **5** (2006), 695 – 718.
10. Gyu Whan Chang and D. Smertnig, *Factorization in the self-idealization of a PID*, *Boll. Unione Mat. Ital.* **IX,6(2)** (2013), 363 – 377.
11. S.T. Chapman, M. Fontana, A. Geroldinger, and B. Olberding (eds.), *Multiplicative Ideal Theory and Factorization Theory*, *Proceedings in Mathematics and Statistics*, vol. 170, Springer, 2016.
12. S.T. Chapman, F. Gotti, and R. Pelayo, *On delta sets and their realizable subsets in Krull monoids with cyclic class groups*, *Colloq. Math.* **137** (2014), 137 – 146.
13. S.T. Chapman, M. Holden, and T. Moore, *Full elasticity in atomic monoids and integral domains*, *Rocky Mt. J. Math.* **36** (2006), 1437 – 1455.
14. S. Colton and N. Kaplan, *The realization problem for delta sets of numerical monoids*, *J. Commut. Algebra*, to appear.
15. S. Frisch, *A construction of integer-valued polynomials with prescribed sets of lengths of factorizations*, *Monatsh. Math.* **171** (2013), 341 – 350.
16. A. Geroldinger, *Additive group theory and non-unique factorizations*, *Combinatorial Number Theory and Additive Group Theory* (A. Geroldinger and I. Ruzsa, eds.), *Advanced Courses in Mathematics CRM Barcelona*, Birkhäuser, 2009, pp. 1 – 86.
17. ———, *Sets of lengths*, *Amer. Math. Monthly* **123** (2016), 960 – 988.
18. A. Geroldinger, D.J. Grynkiewicz, G.J. Schaeffer, and W.A. Schmid, *On the arithmetic of Krull monoids with infinite cyclic class group*, *J. Pure Appl. Algebra* **214** (2010), 2219 – 2250.

19. A. Geroldinger, D.J. Grynkiewicz, and P. Yuan, *On products of k atoms II*, *Mosc. J. Comb. Number Theory* **5** (2015), 73 – 129.
20. A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.
21. A. Geroldinger, W. Hassler, and G. Lettl, *On the arithmetic of strongly primary monoids*, *Semigroup Forum* **75** (2007), 567 – 587.
22. A. Geroldinger, F. Kainrath, and A. Reinhart, *Arithmetic of seminormal weakly Krull monoids and domains*, *J. Algebra* **444** (2015), 201 – 245.
23. A. Geroldinger and M. Roitman, *The arithmetic of one-dimensional local domains*, manuscript.
24. A. Geroldinger and W. A. Schmid, *A characterization of class groups via sets of lengths*, <http://arxiv.org/abs/1503.04679>.
25. ———, *The system of sets of lengths in Krull monoids under set addition*, *Rev. Mat. Iberoam.* **32** (2016), 571 – 588.
26. A. Geroldinger and P. Yuan, *The set of distances in Krull monoids*, *Bull. Lond. Math. Soc.* **44** (2012), 1203 – 1208.
27. A. Geroldinger and Q. Zhong, *A characterization of class groups via sets of lengths II*, *J. Théor. Nombres Bordx.* (2017), (forthcoming).
28. ———, *Long sets of lengths with maximal elasticity*, manuscript.
29. ———, *The set of minimal distances in Krull monoids*, *Acta Arith.* **173** (2016), 97 – 120.
30. F. Halter-Koch, *Ideal Systems. An Introduction to Multiplicative Ideal Theory*, Marcel Dekker, 1998.
31. F. Kainrath, *Factorization in Krull monoids with infinite class group*, *Colloq. Math.* **80** (1999), 23 – 30.
32. J.C. McConnell and J.C. Robson, *Noncommutative Noetherian Rings*, Graduate Studies in Mathematics, vol. 30, American Mathematical Society, 2001.
33. W.A. Schmid, *A realization theorem for sets of lengths*, *J. Number Theory* **129** (2009), 990 – 999.
34. D. Smertnig, *Factorizations in bounded hereditary noetherian prime rings*, <http://arxiv.org/abs/1605.09274>.
35. ———, *Sets of lengths in maximal orders in central simple algebras*, *J. Algebra* **390** (2013), 1 – 43.
36. Q. Zhong, *Sets of minimal distances and characterizations of class groups of Krull monoids*, *Ramanujan J.* (2017), doi:10.1007/s11139-016-9873-2.